

Gamma-ray binaries: pulsars in disguise ?

Guillaume Dubus^{1,2}

¹ Laboratoire Leprince-Ringuet, UMR 7638 CNRS, Ecole Polytechnique, 91128 Palaiseau, France; e-mail: dubus@in2p3.fr

² Institut d'Astrophysique de Paris, UMR 7095 CNRS, Université Pierre & Marie Curie, Paris 6, 98 bis bd Arago, 75014 Paris, France

Draft February 4, 2008

Abstract. LS 5039 and LSI +61°303 are unique amongst high-mass X-ray binaries (HMXB) for their spatially-resolved radio emission and their counterpart at $> \text{GeV}$ gamma-ray energies, canonically attributed to non-thermal particles in an accretion-powered relativistic jet. The only other HMXB known to emit very high energy (VHE) gamma-rays, PSR B1259-63, harbours a non-accreting millisecond pulsar.

The purpose is to investigate whether the interaction of the relativistic wind from a young pulsar with the wind from its stellar companion, as in PSR B1259-63, constitutes a viable scenario to explain the observations of LS 5039 and LSI +61°303. Emission arises from the shocked pulsar wind material, which then flows away to large distances in a comet-shape tail, reproducing on a smaller scale what is observed in isolated, high motion pulsars interacting with the interstellar medium.

The timescales for acceleration and radiation of particles at the shock between the pulsar wind and stellar wind are calculated. Simple expectations for the spectral energy distribution (SED) are derived and are shown to depend on few input parameters. Detailed modelling of the particle evolution is attempted and compared to the observations from radio to TeV energies.

Acceleration at the shock provides high energy electrons that steadily emit synchrotron in X-rays and inverse Compton scatter stellar light to γ -rays. Electrons streaming out of the system emit at IR frequencies and below. The overall aspect of the SEDs is adequately reproduced for standard values of the parameters. The morphology of the radio tail can mimic a microquasar jet. Good agreement is found with the published VLBI map of LS 5039 and predictions are made on the expected change in appearance with orbital phase.

The pulsar wind scenario provides a common, viable framework to interpret the emission from all three γ -ray binaries.

Key words. acceleration of particles — binaries: close — pulsars: general — ISM: jets and outflows — Gamma rays: theory — X-rays: binaries

1. Introduction

X-ray binaries are systems composed of a neutron star or black hole in orbit around a high or low-mass normal star (HMXBs and LMXBs). Non-thermal radio emission has been detected in X-ray binaries and is usually associated with outflows (see Fender, 2006, for a review). In a few objects, radio observations are able to follow the propagation of radio emission up to parsec scales, after initial flaring in the central source. The inferred speeds of the collimated outflow are relativistic with bulk Lorentz factors Γ of a few. These apparently superluminal ejection events have been tentatively associated with transitions from a hard to a soft X-ray spectrum (Fender et al., 2004).

In many more objects, the radio emission has a flat spectrum and is correlated with a hard X-ray spectral state. Usually the radio emission is unresolved but imaging by Very Long Baseline Interferometry (VLBI) of Cyg X-1 in this state has revealed a compact, collimated radio outflow on a scale of a few AU (Stirling et al., 2001). Asymmetries in the surface brightness probably reflect moderate relativistic boosting ($v/c \sim 0.3$). It is as yet unclear whether compact radio emission and super-

luminal ejections can be taken as different manifestations of a unique, underlying ejection mechanism.

Similar types of radio emission (compact or superluminal ejections) are seen in active galactic nuclei (AGN), prompting speculations that phenomenology known to one type of object may have its counterpart in the other. The detection of very high energy γ -rays from the X-ray binary LS 5039 by the HESS collaboration would appear to vindicate this conjecture (Aharonian et al., 2005a). Observations in the 1990s have established that blazars, a subtype of AGN, emit high energy (HE, GeV) and very high energy (VHE, TeV) γ -rays (Hoffman et al., 1999). Blazars have their jet aligned close to the line-of-sight, resulting in a very strong relativistic boosting of the jet non-thermal emission. The same configuration could be found in an X-ray binary and this is perhaps the case in LS 5039.

LS 5039 is composed of a O6.5V star and an unidentified compact object in a 3.9 day orbit (Motch et al., 1997; Clark et al., 2001; McSwain & Gies, 2002; McSwain et al., 2004; Casares et al., 2005b). Much of the interest in this object stems from its tentative association with the EGRET source

3EG J1824-1514 and from the fact that its radio emission was shown to be extended on a milliarcsecond scale (Paredes et al., 2000). By analogy with blazars, it is tempting to attribute the radio and γ -ray emission to high energy particles accelerated in a jet. LS 5039 is very similar to LSI +61°303 an X-ray binary with a B0Ve star in a 26.5 day orbit whose association with a high energy source dates back to COS-B (Gregory & Taylor, 1978; Hutchings & Crampton, 1981; Casares et al., 2005a). The associations rely purely on positional coincidence and this is fairly poor, the EGRET 99% confidence levels being roughly at 0.5° for LS 5039 and 0.2° for LSI +61°303 (3EG J0241+6103). The HESS detection of VHE emission coincident to within an arcmin with the position of LS 5039 identifies this X-ray binary as the source of γ -rays. By extension, it places the EGRET tentative identification on a firmer footing.

VHE γ -ray emission is at least evidence that particles can be accelerated to very high energies in X-ray binaries. However, is it necessarily associated with blazar-like model, involving accretion of material, ejection and non-thermal acceleration in a relativistic jet? Maraschi & Treves (1981) pointed out early on that the spindown of a young, rapidly-rotating pulsar could power the γ -ray emission of LSI +61°303. The relativistic pulsar wind would be confined by the stellar wind from the massive companion. Particles accelerated at the termination shock then produce the non-thermal emission.

There is at least one system where such a scenario is undoubtedly at work: PSR B1259-63 (=SS 2883), a 47.7 millisecond (ms) radio pulsar in a 3.4 yr orbit around a B2Ve star (Johnston et al., 1994; Manchester et al., 1995). Variable γ -ray emission was detected by HESS during periastron passage (Aharonian et al., 2005b). In LS 5039 and LSI +61°303, the discovery of resolved radio emission was interpreted as the signature of a relativistic jet and recently, the attention seems to have focused on accretion/ejection scenarios (but see Leahy 2004; Martocchia et al. 2005). The pulsar scenario has become unwonted. Yet, pulsars interacting with their surrounding medium can also lead to large scale emission, the bow shock nebula of high velocity ms pulsars even taking a comet tail appearance (e.g. Gaensler, 2005). The resolved ‘jets’ of LS 5039 and LSI +61°303 might be small-scale analogues, but with the trail originating from the interaction with a companion wind rather than with the ISM.

The similarities between LS 5039 and LSI +61°303 have long been noticed: resolved radio emission, hard steady X-ray fluxes and EGRET emission in the GeV range at a level $\sim 10^{35}$ erg s⁻¹. LSI +61°303, located in the Northern hemisphere and inaccessible to HESS, has yet to be detected at TeV energies. While PSR B1259-63 shares the property of being a TeV emitter with LS 5039, LSI +61°303 and PSR B1259-63 both have similar periodic radio outbursts that may be linked to both of them having a Be type companion. PSR B1259-63 has not been detected at GeV energies by EGRET (including at periastron passage, Tavani et al., 1996), perhaps because its long orbit only takes it close to the star for periods of time that are too limited for the sensitivity of the instrument.

At closest, PSR B1259-63 is about 0.7 AU away from its companion, equivalent to the apastron orbital separation in LSI +61°303. The periastron separation of 0.1 AU in

Table 1. Orbital parameters adopted for the γ -ray binaries.

System	P_{orb} (d)	e	M_2 (M_{\odot})	d_S (AU)	R_{\star} (R_{\odot})	T_{\star} (10^4 K)
LS 5039	3.9	0.35	23	0.1-0.2	9.3	3.9
LSI +61°303	26.5	0.72	12	0.1-0.7	10	2.2
PSR B1259-63	1237	0.87	10	0.7-10	10	2.7

LSI +61°303 is itself comparable to the orbit in the more compact LS 5039 (Table 1). There is a scaling between the three systems that, should they be governed by the same underlying processes, might lead to interesting comparisons. Indeed, their spectral energy distributions are similar in shape and luminosity (Fig. 1), suggesting the same physics might be at work in these three known examples of “ γ -ray binaries”.

This work purports to investigate the pulsar wind scenario in light of the latest observations of these binaries. Reviewing current evidence, it is argued in §2 that there is as yet no reason to discard this possibility in LS 5039 and LSI +61°303. The scenario is then developed in the following sections. The basic assumptions are described in §3, showing how the physical quantities scale with the parameters of the model. Simple expectations for the spectral energy distribution are derived in §4. A numerical model of the emission is elaborated in §5 and applied to the γ -ray binaries in §6. It is shown that the radio emission from the shocked pulsar wind material can look like that of a compact jet emanating from an accreting microquasar. The conclusion set out in §7 is that the pulsar wind scenario provides a plausible, coherent and persuasive framework to interpret the emission from these γ -ray binaries.

2. Accretion or rotation-powered ?

Several observations could securely tilt the balance towards the ‘black hole jet’, accretion-powered scenario or the ‘pulsar wind’, rotation-powered scenario in LS 5039 and LSI +61°303 without detailed modelling. These are examined here in turn.

2.1. Is the compact object a black hole ?

A compact object mass greater than 3 M_{\odot} would rule out the pulsar wind scenario. Constraints have been set in both LS 5039 and LSI +61°303 by measuring the radial velocity of the stellar companion. This enables the measure of the period, mass function, eccentricity and periastron angle. The mass function provides a strict lower limit to the mass of the compact object. In high-mass X-ray binaries the mass function is too small to provide a useful constraint: $f(M) = 0.053 M_{\odot}$ in LS 5039, $f(M) = 0.011 M_{\odot}$ in LSI +61°303 (Casares et al. 2005a,b who have the best available datasets).

A constraint on the compact object mass then requires additional knowledge on the mass of the stellar companion and the orbit inclination. Atmosphere fitting of the spectrum leads to an estimate for the mass (and radius) of the massive star. Accuracy requires detailed stellar models and is inherently difficult due to the high mass loss rates of early-type stars. Lines can originate at different depths and veloci-

ties in the strong (variable) stellar wind, leading to confusing measurements of radial velocities and difficulties in spectral modelling (see e.g. discussion in Bahcall, 1978). Parameters thereby derived have higher, probably irreducible, systematics than the quoted (statistical) errors might suggest (e.g. Table 2 of Casares et al., 2005b). In fact, the orbital period of LSI +61°303 (26.5 days) cannot be determined from optical spectroscopy alone with high confidence and radial velocity studies use as an *a priori* the well-determined radio outburst period (Hutchings & Crampton, 1981; Gregory, 2002; Casares et al., 2005a).

The parameter space derived by Casares et al. (2005a,b) using these techniques allows for both a neutron star or a black hole in LS 5039 and LSI +61°303. The radial velocity amplitude implies a low orbit inclination if the compact object is massive ($i \approx 30^\circ$ for $M > 3M_\odot$ in both LS 5039 and LSI +61°303), a good omen for a model with a relativistic jet pointed close to the line-of-sight. On the other hand, a $1.4 M_\odot$ neutron star implies in both objects $i \approx 60^\circ$. In LS 5039, the lack of X-ray eclipses (assuming point-like emission) by a companion of $\approx 9 R_\odot$ requires $i \lesssim 65^\circ$, consistent with the above. Line broadening of the lines gives an upper limit on their rotation speeds which, since it must be lower than breakup speed, can be used to set a lower limit on the inclination for aligned spin/orbital axis. Casares et al. (2005a,b) find $i > 10^\circ$ in both systems, again consistent with both a black hole or neutron star. Since the value of $\sin i$ is more likely to be high than low for a random distribution of inclinations, a neutron star would seem to be favoured over a black hole.

The binary orbit of LS 5039 is quite compact for a high-mass X-ray binary ($P_{\text{orb}} \approx 3.9$ days) and is only mildly eccentric ($e \approx 0.35$). By comparison, LSI +61°303 has $P_{\text{orb}} \approx 26.5$ days and $e \approx 0.7$. Orbital circularization and synchronization might be relatively advanced in LS 5039. This has prompted Casares et al. (2005b) to use their measurement of rotational broadening to constrain the orbit in LS 5039 by assuming that the massive star is corotating at the angular orbital velocity at periastron. The compact object mass then becomes $2.7\text{--}5.0 M_\odot$ with $i = 24.9 \pm 2.8^\circ$, probably optimistic because of the possible systematics, but in practice ruling out a neutron star.

This conclusion is not robust. Observations of the stellar companions in other high-mass X-ray binaries with similar periods do not systematically show corotation (Conti, 1978). Although synchronization is undoubtedly fast by astronomical standards (some detached early-type binaries are observed to be circularized), short-lived wind-fed HMXBs do not have ages much greater than their typical synchronization timescale. These timescales are difficult to calculate precisely: in massive stars, they depend on excited modes in the radiative envelope (Zahn, 1977). The age of the system can only be guessed at. Rotation-powered emission would probably imply an energetic pulsar with a short spindown timescale. Synchronization would then not have happened.

In both LSI +61°303 and LS 5039, optical spectroscopy alone is unable to decide between a neutron star and black hole compact object. This work will assume a $1.4 M_\odot$ neutron star.

2.2. Is there accretion?

Any sign of accretion on the compact object would dismiss the pulsar wind scenario. Accretion would have to be wind-fed: Roche lobe overflow as occurs in Cyg X-1 and some other HMXBs would entail much larger X-ray luminosities. The orbits and sizes of the massive stars in LS 5039 and LSI +61°303 support this conclusion (Casares et al., 2005a,b). The rate at which mass is accreted from the stellar wind can be estimated. The Bondi capture radius is $r_a = 2GM_c/v_w^2$, where v_w is the wind speed (about 2000 km s^{-1}) and M_c is the mass of the compact object. The Bondi mass accretion rate is then $\dot{M} \approx \dot{M}_w/(2r_a/d_s)^2$ where \dot{M}_w is the stellar wind mass loss rate (about $10^{-7} M_\odot \text{ yr}^{-1}$) and d_s is the orbital separation. Using the LS 5039 orbital and stellar parameters in Table 1 gives between $0.6\text{--}20 \times 10^{14} \text{ g s}^{-1}$ along the orbit in LS 5039, with a time-average of $5 \times 10^{14} \text{ g s}^{-1}$. With 10% radiative efficiency, this is too close to the sustained GeV emission at a level $\sim 10^{35} \text{ erg s}^{-1}$. In addition, although the X-ray luminosity is comparable to those of wind-fed X-ray binaries, the lightcurve does not show evidence for the large expected associated orbital variability (Reig et al., 2003; Bosch-Ramon et al., 2005). The slow, dense Be equatorial wind in LSI +61°303 allows for higher accretion rates, in better agreement with the total luminosity and the factor ~ 10 variations observed in the 0.5-2 keV band lightcurve (less above, Harrison et al. 2000; Martí & Paredes 1995, Paredes et al. 1997).

There are no signs of the large excursions in luminosity or spectra typical of (hard or soft) X-ray transients. The X-ray spectra are reminiscent of the low/hard X-ray state of LMXBs. There are some radio/X-ray fluctuations on timescales \sim hour in both objects (Harrison et al., 2000; Bosch-Ramon et al., 2005); but strong, fast variability on timescales of \sim second, expected from a low/hard state LMXB or accreting HMXB (van der Klis, 2006), has not been detected (Ribó et al., 1999; Harrison et al., 2000, although the low flux of the sources should be noted). The Thomson opacity is low in the stellar wind, hence variability might be smoothed out by scattering in a dense corona. There is no sign of such a corona in X-ray spectra. Indeed, the apparent lack of spectral cutoffs around either 30 keV or 100 keV (see Fig. 1) puts them apart from both accreting HMXBs and low state LMXBs. *RXTE* spectra of LS 5039 did show a strong, broad Fe line (Ribó et al., 1999), but that could not be confirmed by other X-ray missions (Martocchia et al., 2005). LS 5039 lies in the Galactic Plane and the Fe line is very likely due to Ridge emission sampled by the large *RXTE* non-imaging aperture (Bosch-Ramon et al., 2005).

The radio and hard X-ray emission would be compatible with most of the emission coming from a non-thermal distribution of particles in a relativistic jet (Romero et al., 2005; Paredes et al., 2006; Dermer & Böttcher, 2006). Gamma-rays would then be up-scattered photons. The puzzle is how such a jet could be fed by accretion without any of the usual signatures, most notably variability. In the γ -ray binaries, the broad-band SEDs are stable even on timescales of years.

The simplest hypothesis is that, as in PSR B1259-63, accretion does *not* occur in LS 5039 and LSI +61°303. If the

infall of material is stopped by pressure from a relativistic pulsar wind, this will produce very little emission and variability. Instead, the emission will be due to the shocked pulsar wind. As in the relativistic jet scenario, a non-thermal distribution of particles provides the necessary ingredient to explain the SED. The spindown of the pulsar provides a stable energy source on human timescales. The termination shock of the contained pulsar wind provides the conditions for particle acceleration. Only very mild variations are expected, due to fluctuations of the environment (stellar wind), consistent with observations.

2.3. Is the radio emission a microquasar jet?

The detection of relativistic motion in the radio emission, like that of the microquasar GRS 1915+105, would secure the jet model. The radio morphology of LS 5039 and LSI +61°303 looks more like the compact jets seen in the low/hard X-ray state of low-mass X-ray binaries or Cyg X-1, with the significant difference that their radio spectral indices are steeper than the usual flat indices measured in the self-absorbed jets. The indices ($F_\nu \sim \nu^{-0.5}$) rather hint at optically thin synchrotron emission from a $dN \propto E^{-2}dE$ power-law distribution of electrons.

The period of the radio outbursts of LSI +61°303 is very stable and is used as a clock for the orbital motion (Gregory, 2002). The radio luminosity of LS 5039 has been steady at the same level for years (Martí et al., 1998a). The radio brightness asymmetry of LS 5039 and LSI +61°303 has been used to place constraints on (de)boosting of the putative relativistic jet. This yields a mild bulk flow velocity of 0.2-0.3c (Paredes et al., 2000; Massi et al., 2004). There is no evidence in either system for highly relativistic bulk motion directed more or less towards the line-of-sight; the main argument favouring a jet interpretation is based on the morphology of the radio emission.

As mentioned in §1, fast-moving young and old millisecond pulsars leave trails of X-ray, optical and radio emission that can reach parsec lengths, and whose properties scale with the distance where the wind terminates due to containment by the surrounding interstellar medium (Wang et al., 1993; Arons & Tavani, 1993). Even the collision of standard stellar winds can produce detectable large-scale, jet-looking, radio emission (e.g. Dougherty et al., 2003). The radio emission of LS 5039 and LSI +61°303, seems to reach scales of up to several 100 AU (Paredes et al., 2002; Massi et al., 2004), double-sided in the case of LS 5039. Energetically speaking, this emission can easily be a pulsar nebula. However, isolated ms pulsars have time to develop long (one-sided) nebula away from their direction of proper motion. In the γ -ray binaries, the direction of such a nebula would be affected on short timescales by orbital motion. Radio observations of LSI +61°303 have shown curved radio emission on a scale of 50 mas, interpreted as due to precession of a jet as in SS 433 (Massi et al., 2004). Could this morphology instead be a result of the pulsar orbital motion?

In this work, it will be argued that the resolved radio emission in LS 5039 and LSI +61°303 is *not* due to a relativistic jet akin to those of AGN or microquasars (and implying accre-

tion), but arises from shocked pulsar wind material outflowing from the binary.

2.4. Can the compact object be a pulsar?

The X-ray and radio properties of LS 5039 and LSI +61°303 sets them apart from other X-ray binaries, with no conclusive signs of accretion going on. At the same time, the luminosity, hard X-ray power law extending to high energies, weak variability and optically thin radio emission are typical for emission from young pulsars with a high spindown power (e.g. Gaensler & Slane, 2006). Examining the pulsar wind hypothesis therefore seems justified.

In order to avoid disc or magnetospheric accretion, the relativistic wind of the pulsar must be able to quench the infall of stellar matter attracted to it (Illarionov & Sunyaev, 1975). If all of the pulsar spindown power \dot{E} is transferred to this relativistic wind, then writing the pressures gives $\dot{E} > 4\dot{M}v_w c$ where \dot{M} is the Bondi rate calculated above. Writing the spindown power as a function of the pulsar magnetic field B and period P , then $P \lesssim 230B_{12}^{1/2}\dot{M}_{15}^{-1/4}$ ms, with $\dot{M}_{15} = \dot{M}/10^{15} \text{ g s}^{-1}$. A young, ms pulsar would push out any settling accretion flow.

The young age is not problematic. Radio searches have not conclusively identified an association with a supernova remnant (Frail et al., 1987; Martí et al., 1998a; Ribó et al., 2002); but neither have they been able to do so for PSR B1259-63, which has a measured spindown age of $3 \cdot 10^5$ years (Johnston et al., 1992). Ribó et al. (2002) put an upper limit of about a Myr to the age of LS 5039 by tracing back the proper motion of the system to the plane. Nitrogen enrichment in the companion's atmosphere also suggests a young age in LS 5039 (McSwain et al. 2004; rotational-mixing could also explain the enrichment, Casares et al. 2005b).

Detecting the radio pulse would nail down this scenario. Unfortunately, free-free absorption in the dense stellar winds of LS 5039 and LSI +61°303 would suppress the signal. Using the Rosseland mean opacity, the radial optical depth at 1 GHz from a point at a distance d from the star is $\tau \approx 3 \cdot 10^4 \dot{M}_{w,7}^2 v_{w,2000}^{-2} T_{4}^{-3/2} \nu_{\text{GHz}}^{-2} d_{0.1}^{-3}$ for a $10^{-7} M_\odot \text{ yr}^{-1}$ coasting wind with a speed 2000 km s⁻¹ and temperature 10⁴ K; d is taken to be 0.1 AU, about the periastron orbital separation in LS 5039 and LSI +61°303. High opacities are also found in LSI +61°303 (Taylor & Gregory, 1982). In PSR B1259-63, the suppression of pulsed emission is observed close to periastron (Johnston et al., 1992; Melatos et al., 1995).

Pulsed emission in the X-ray/GeV band would serve just as well. Such a detection does not seem realistic without prior knowledge on the pulse period derivatives given the sensitivities achievable even by *GLAST*. In addition, the pulsed GeV signal in LS 5039 may be smoothed out by emission from the e^+e^- cascade initiated by absorption of TeV γ -rays on stellar photons (Dubus, 2006).

As the pulsar rotation gradually decreases, the γ -ray binaries would become standard accretion-powered high mass X-ray binaries, perhaps brighter in the X-ray sky but probably without significant emission at γ -ray energies. That three examples of such short-lived phases may have been found

perhaps implies a large population of descendant HMXBs of which some might be the enshrouded hard X-ray sources discovered by *INTEGRAL*. Some of the unidentified EGRET sources located in the Galactic Plane may also turn out to be pulsars in binaries. Massive X-ray binary population synthesis calculations by Meurs & van den Heuvel (1989) predict about 30 binaries in the brief young pulsar stage in the Galaxy, consistent with three examples being found within 3 kpc.

The conclusion is that LS 5039 and LSI +61°303 could very well harbour a young pulsar with a strong relativistic wind like that of PSR B1259-63. A model based on this scenario is explored in the next sections.

3. Scaling relationships in the binary plerion

The scenario, identical to that proposed for PSR B1259-63, is the following: a young pulsar with a spindown rate of order 10^{36} erg s⁻¹ generates a strong relativistic wind beyond the light cylinder (Rees & Gunn, 1974). The pulsar wind is assumed to be radial, isotropic and composed of principally mono-energetic electrons/positrons with a Lorentz factor γ_p . All the spindown power is transferred to the leptons, with a small fraction going to the magnetic field. The ratio of magnetic to kinetic energy σ is a parameter of the model.

The pulsar wind is contained by the stellar wind (in a plerion like the Crab, the container is the supernova remnant material). A collisionless shock forms beyond which the leptons from the pulsar wind are accelerated and isotropized. The shocked pulsar wind material is separated from the stellar wind by a contact discontinuity across which mixing might occur (neglected here). The shock has a ‘bow’ or ‘comet’ shape with a tail extending away from the stellar companion. The radio emission occurs in this nebula. Studying such a complex interaction is simplified by noting everything scales with the standoff distance (see e.g. Bucciantini et al., 2005). The standoff distance R_s is the radial distance from the pulsar to the star at which the ram pressures equilibrate (§3.2).

Most emission occurs in the vicinity of R_s , whose value can be calculated given the orbital parameters, and the stellar wind density and velocity. Straightforward estimates indicate that the physical conditions at the shock are set by only three quantities: the shock distance R_s from the pulsar, the orbital separation d_s and the combination $\sigma\dot{E}$, which describes the pulsar wind energy and magnetization. The overall properties of the resulting spectral energy distribution are then deduced in the next section (§4). A quantitative description including emission from the nebula is given in §5-6.

3.1. The pulsar wind

The pulsar wind is described by the Lorentz factor γ_p of the electrons/positrons, the spindown power \dot{E} and the ratio of magnetic to kinetic energy σ . The (proper frame) density n_1 and (observer frame) magnetic field B_1 in the pulsar wind upstream of the shock are then derived from (Kennel & Coroniti,

1984b)

$$\sigma = \frac{B_1^2}{4\pi n_1 \gamma_p^2 m_e c^2} \quad (1)$$

$$\frac{\dot{E}}{4\pi R_s^2 c} = \frac{B_1^2}{4\pi} \left(\frac{1 + \sigma}{\sigma} \right) \quad (2)$$

Most of the energy could actually be carried by ions, which are probably a necessary ingredient to accelerate the leptons at the termination shock (Arons & Tavani, 1994). Here, the kinetic energy of the wind is assumed to be only in the electrons/positrons.

In PSR B1259-63 \dot{E} is $8 \cdot 10^{35}$ erg s⁻¹ and the pulsar magnetic field B is about $3 \cdot 10^{11}$ G (corresponding to a spin period $P = 47.7$ ms and period derivative $\dot{P} = 2.3 \cdot 10^{-15}$). Thereafter, \dot{E} will be expected to be of order 10^{36} erg s⁻¹. Models of the Crab plerion yield values of order 10^{-3} for σ and γ_p in the $10^5 - 10^6$ range (Rees & Gunn, 1974; Kennel & Coroniti, 1984a). Explaining the implied conversion of electromagnetic energy (close to the pulsar) to kinetic energy (far from the pulsar) is a key question of pulsar physics. Values for σ could be higher if the shock occurs closer to the pulsar than in Crab, as proposed for PSR B1259-63 by Tavani & Arons (1997); in any case σ will be expected smaller than 1.

3.2. The standoff distance

The standoff distance point R_s of the pulsar wind is given by

$$\frac{\dot{E}}{4\pi R_s^2 c} = \rho_w (\mathbf{v}_w - \mathbf{v}_p)^2 \quad (3)$$

where \mathbf{v}_p is the pulsar radial orbital speed calculated from the binary parameters, ρ_w is the stellar wind density and \mathbf{v}_w the stellar wind speed at the standoff distance. An estimate of R_s can be obtained by balancing the pressures of the two flows, assuming a stellar wind coasting at a speed v_∞ and a negligible orbital speed:

$$R_s \approx \frac{d_s}{1 + (P_w/\dot{E})^{1/2}} \quad (4)$$

where $P_w/c = \dot{M}_w v_\infty$ is the stellar wind momentum flux with \dot{M}_w the stellar mass loss rate, supposed isotropic here, and d_s is the orbital separation. To fix ideas, the radiatively-driven wind of LS 5039 has $\dot{M}_w \approx 10^{-7} M_\odot$ yr⁻¹ and $v_\infty \approx 2000$ km s⁻¹ which gives $P_w \approx 5 \cdot 10^{37}$ erg s⁻¹ (McSwain et al., 2004). When the stellar wind is more powerful than the pulsar wind, in the sense that $P_w \gg \dot{E}$, then the standoff point can be written as

$$R_s \approx 1.5 \cdot 10^{11} \text{ cm } d_{0.1} P_{w,38}^{-1/2} \dot{E}_{36}^{1/2} \quad (5)$$

where $d_{0.1} = d_s/0.1$ AU is the orbital separation (0.1 AU at periastron in LS 5039 and LSI +61°303), $\dot{E}_{36} = \dot{E}/10^{36}$ erg s⁻¹ and $P_{w,38} = \dot{M}_w v_\infty c/10^{38}$ erg s⁻¹. The standoff distance stays at about a tenth of the binary separation throughout the orbit. Writing down explicitly the dependence on the stellar wind (Eq. 5) is easy using v_∞ and P_w . Eq. 4 is used for the slow equatorial winds of Be stars (§6). In the following, it proved more convenient to explicitly keep R_s as a parameter, derived

under the plausible assumptions described above. Some of the uncertainties in the geometry of the shock region or the stellar wind parameters (particularly in the wind acceleration region) are thereby hidden in R_s .

3.3. Downstream flow at standoff

Downstream values for the magnetic field and particle density are calculated using perpendicular MHD shock jump conditions at R_s (Kennel & Coroniti, 1984a). The post-shock magnetic field is (for $\sigma \ll 1$)

$$B = 3(1 - 4\sigma) \left(\frac{\dot{E}/c}{R_s^2} \frac{\sigma}{1 + \sigma} \right)^{1/2} \approx 5 (\dot{E}_{36} \sigma_3)^{1/2} R_{11}^{-1} \text{ G} \quad (6)$$

where $\sigma_3 = \sigma/10^{-3}$ and $R_{11} = R_s/10^{11} \text{ cm}$.

The downstream particles are assumed to be accelerated to a canonical $dn_\gamma \propto \gamma^{-2} d\gamma$ power law from a minimum energy γ_{\min} to a maximum energy γ_{\max} . For a given γ_{\max} , the normalisation and lower limit γ_{\min} of the post-shock particle distribution are found by matching to the downstream particle and energy densities. For a γ^{-2} power law and $\sigma \ll 1$, γ_{\min} is given by (Kennel & Coroniti, 1984b)

$$\gamma_{\min} \ln \left(\frac{\gamma_{\max}}{\gamma_{\min}} \right) \approx \frac{\gamma_p}{\sqrt{2}} \quad (7)$$

For a post-shock distribution covering 2-3 decades in energy, $\gamma_{\min} \approx 0.1\gamma_p$.

3.4. Timescales at standoff

Particles are assumed to be accelerated efficiently up to the maximum energy γ_{\max} above which the timescale at which particles radiate their energy becomes shorter than the acceleration timescale. The gyroradius is used to give an estimate of the acceleration timescale to a Lorentz factor γ

$$t_{\text{acc}} \approx \frac{\gamma m_e c}{eB} \approx 0.01 \gamma_6 (\dot{E}_{36} \sigma_3)^{-1/2} R_{11} \text{ seconds} \quad (8)$$

using the downstream magnetic field (Eq. 6) and writing $\gamma_6 = \gamma/10^6$. For ion dominated shocks (Arons & Tavani, 1994), the relevant timescale is given by the ion gyroradius and the maximum electron energy is $\sim \gamma_p m_p/m_e$ (γ_p is then the upstream ion Lorentz factor). The maximum γ reached must have a gyroradius smaller than the characteristic scale of the shock R_s , implying $\gamma < 3 \cdot 10^8 (\dot{E}_{36} \sigma_3)^{1/2}$. Particles do not reach such high γ as they then lose energy to radiation more rapidly than they can be accelerated.

The synchrotron timescale is

$$t_{\text{sync}} = \frac{9 m_e^3 c^5}{4 e^4 \gamma} \frac{1}{B^2} \approx 30 \gamma_6^{-1} (\dot{E}_{36} \sigma_3)^{-1} R_{11}^2 \text{ seconds} \quad (9)$$

so the maximum γ such that $t_{\text{sync}} > t_{\text{acc}}$ is

$$\gamma_{\max} \approx 5 \cdot 10^7 (\dot{E}_{36} \sigma_3)^{-1/4} R_{11}^{1/2} \quad (10)$$

which, in principle, could result in a maximum photon energy of 25 TeV. Photons up to energies $\gtrsim 4 \text{ TeV}$ are detected by HESS (Aharonian et al., 2005a), requiring electrons

with Lorentz factors high enough that $\gamma_{\max} m_e c^2 \gtrsim 4 \text{ TeV}$, i.e. $\gamma_{\max} \gtrsim 10^7$. This is below the limit from synchrotron losses (Eq. 10) for all plausible values of the pulsar σ -parameter.

The inverse Compton timescale in the Klein-Nishina regime $\gamma h\nu \gg m_e c^2$ is (Blumenthal & Gould, 1970)

$$t_{\text{KN}} \approx 5.2 \gamma_6 \left(\frac{3 \text{ eV}}{kT_\star} \right)^2 \left(\frac{d_s}{R_\star} \right)^2 \left[\ln \left(\frac{4\gamma kT_\star}{m_e c^2} \right) - 1.98 \right]^{-1} \text{ seconds} \quad (11)$$

where the stellar (blackbody) radiation density has been lowered by geometric dilution and R_s is assumed to be small compared to the orbital separation d_s . The stellar temperature T_\star varies between $2\text{--}4 \cdot 10^4 \text{ K}$ with the stellar radius $R_\star \approx 10 R_\odot$ for LS 5039, LSI +61°303 and PSR B1259-63 so that

$$t_{\text{KN}} = 20 \gamma_6 d_{0.1}^2 [\ln \gamma_6 + 1.3]^{-1} T_{\star,4}^{-2} R_{\star,10}^{-2} \text{ seconds} \quad (12)$$

with $T_{\star,4} = T_\star/40,000 \text{ K}$ and $R_{\star,10} = R_\star/10 R_\odot$. In the Thomson limit $\gamma h\nu \ll m_e c^2$ then

$$t_{\text{T}} = \frac{9 m_e^3 c^5}{4 e^4 \gamma} \frac{1}{8\pi U_\star} \approx 40 \gamma_3^{-1} d_{0.1}^2 T_{\star,4}^{-4} R_{\star,10}^{-2} \text{ seconds} \quad (13)$$

where $U_\star = (\sigma_B T_\star^4/c)(R_\star/d_s)^2$ is the stellar photon density. Most of the stellar photons have an energy of $2.7 kT_\star \approx 9 \text{ eV}$ so that the turning point between Thomson and Klein-Nishina regime is for $\gamma_{\text{KN}} \approx 511 \text{ keV}/9 \text{ eV} \approx 6 \cdot 10^4$. The interaction timescale is then at a minimum of $\approx 7 \text{ s}$ (see Fig. 2).

The timescale for particles in the pulsar wind to reach the standoff point is

$$t_w \approx R_s/c \approx 3 R_{11} \text{ seconds} \quad (14)$$

which is only slightly shorter than the minimum inverse Compton timescale derived above. This is important since if the pulsar wind is composed of mono-energetic particles with $\gamma_p \approx 6 \cdot 10^4$, then a fraction of the spindown power can be radiated by bulk inverse Compton interactions in γ -rays before reaching the termination point (Compton drag). There may then not even be a shock. (The upstream magnetic field is frozen in the MHD flow so upstream particles do not radiate synchrotron.) The resulting γ -ray spectrum would display line-like emission with a strong orbital phase dependence (Bogovalov & Aharonian, 2000; Ball & Kirk, 2000). This possibility appears excluded by the power-law HESS spectra extending to a few TeV in LS 5039 and PSR B1259-63 (Aharonian et al., 2005a,b). Compton losses are weaker in LSI +61°303 and PSR B1259-63 (smaller T_\star and larger separations) than in LS 5039.

Downstream of the shock, the inverse Compton timescale is greater than the synchrotron timescale when $\gamma < \gamma_S$ where

$$\gamma_S \approx 1.2 \cdot 10^6 (R_{11}/d_{0.1}) (\dot{E}_{36} \sigma_3)^{-1/2} T_{\star,4} R_{\star,10} \quad (15)$$

using Eq. 9 and 12. The maximum electron energy γ_{\max} is indeed set by synchrotron losses (Eq. 10), and not by inverse Compton losses. Perhaps counter-intuitively, high energy particles lose their energy mostly through synchrotron radiation, although the radiation density at R_s from the luminous stellar companion is much larger than the magnetic field energy density ($U_\star/U_B \approx 10^4 (R_{11}/d_{0.1})^2 (\dot{E}_{36} \sigma_3)^{-1}$). The difference in densities is compensated by the reduction of the inverse

Compton cross-section in the Klein-Nishina regime. Note that γ_S is independent of the orbital separation when the stellar wind dominates ($R_s \propto d_s$, see Eq. 5).

When $\sigma \ll 1$, the post-shock material is found from the jump conditions to initially flow away at $v \approx c/3$ (Kennel & Coroniti, 1984a) so the escape timescale is of order

$$t_{\text{esc}} \approx \frac{R_s}{c/3} \approx 10 R_{11} \text{ seconds} \quad (16)$$

Here, $\gamma_{\min} \approx 0.1\gamma_p > \gamma_{\text{KN}}$ so particles radiate in the Klein-Nishina regime up to γ_S . Their radiative timescale $t_{\text{IC}} \approx 7-20$ s is comparable to their escape timescale. Particles with $\gamma > \gamma_S$ lose energy to synchrotron radiation, on a timescale $t_{\text{sync}} \lesssim 30$ s that is, again, comparable to their escape timescale. The synchrotron radiative efficiency increases with decreasing R_s since $t_{\text{sync}}/t_{\text{esc}} \sim R_s$.

4. Expected high energy spectrum

The timescales given above are sufficient to get an approximate idea of the expected high energy emission from the pulsar wind shock region. The aim is to provide a simple framework to understand the results from the more complex modelling presented in §5-6.

4.1. Spectral shape

Since the radiation timescales are comparable to or (much) smaller than the flow timescale, the situation can be approximated to steady-state injection of power-law electrons interacting in the Klein-Nishina regime with an isotropic bath of quasi mono-energetic photons. Moderski et al. (2005) have studied exactly this case, taking into account synchrotron emission in a uniform magnetic field. The overall characteristics of the spectrum emitted in the vicinity of the pulsar can be deduced from their results and the calculations above. If particle cooling is not fast enough, then particles will travel to regions of lower energy densities, modifying the spectral energy distribution from that presented here. The steady state assumption is reasonable for the high energy spectrum (§5).

Moderski et al. (2005) show that the steady-state electron distribution displays three regimes: (a) at low energies, interactions in the Thomson limit steepen, as usual, the distribution by 1 in index; (b) at high energies, synchrotron emission also steepens the distribution by 1; (c) at intermediate energies, where inverse Compton interactions in the Klein-Nishina regime dominate, the decrease in cross-section is such that the distribution index is unchanged or hardened at most by $\lesssim 0.5$. The hardening decreases with the ratio U_\star/U_B . The change between regimes (a) and (b) occurs at the transition to the Klein-Nishina regime γ_{KN} . Case (a) does not happen here since we initially have $\gamma_{\min} > \gamma_{\text{KN}}$. The change between regimes (b) and (c) occurs at γ_S where synchrotron losses start to dominate. A $dN \sim \gamma^{-2} d\gamma$ injection spectrum will therefore lead to a steady-state electron distribution going from $dN \sim \gamma^{-1.5} d\gamma$ to γ^{-3} above γ_S .

Using the downstream magnetic field (Eq. 6), the characteristic synchrotron frequency for an electron with an energy $\gamma m_e c^2$ is

$$h\nu \approx 100 \gamma_6^2 (\dot{E}_{36} \sigma_3)^{1/2} R_{11}^{-1} \text{ keV} \quad (17)$$

The three frequencies relevant to the spectral shape, corresponding to γ_{\min} , γ_S and γ_{\max} are (using $\gamma_{\min} \approx 0.1\gamma_p$ and Eqs. 10,15):

$$h\nu_{\text{sync,min}} \approx 1 \gamma_{p,6}^2 (\dot{E}_{36} \sigma_3)^{1/2} R_{11}^{-1} \text{ keV} \quad (18)$$

$$h\nu_{\text{sync,S}} \approx 150 (\dot{E}_{36} \sigma_3)^{-1/2} (T_{\star,4} R_{\star,10})^2 (R_{11}/d_{0.1}^2) \text{ keV} \quad (19)$$

$$h\nu_{\text{sync,max}} \approx 250 \text{ MeV} \quad (20)$$

The expected synchrotron spectrum (Moderski et al., 2005) is a hard power law $\nu F_\nu \sim \nu^{0.7}$ from about 1 keV initially (γ_{\min}) up to a break energy ≈ 100 keV (γ_S). The spectrum then flattens, with $\nu F_\nu \sim \text{constant}$ up to ≈ 250 MeV (γ_{\max}). Since R_s/d_s is almost constant (Eq. 5), the break frequency changes as d_s^{-1} along the orbit because of the variation of the magnetic field at the standoff point. The maximum synchrotron frequency is independent of all parameters.

Inverse Compton interactions occur in the Klein-Nishina regime so that the characteristic frequencies are given by $h\nu \approx \gamma m_e c^2$

$$h\nu_{\text{comp,min}} \approx 50 \gamma_{p,6} \text{ GeV} \quad (21)$$

$$h\nu_{\text{comp,S}} \approx 0.6 (\dot{E}_{36} \sigma_3)^{-1/2} (T_{\star,4} R_{\star,10}) (R_{11}/d_{0.1}) \text{ TeV} \quad (22)$$

$$h\nu_{\text{comp,max}} \approx 25 (\dot{E}_{36} \sigma_3)^{-1/4} R_{11}^{1/2} \text{ TeV} \quad (23)$$

Moderski et al. (2005) find the inverse Compton spectrum for a γ^{-2} injection is constant or rising slightly in νF_ν up to $\nu_{\text{comp,S}}$ (their Fig. 6). Above an energy $\gamma_S m_e c^2$, the electron energy distribution steepens to γ^{-3} . The spectral index steepens by ≈ 1.5 (their Eq. 31) and $\nu F_\nu \sim \nu^{-1.5}$ is expected above 0.6 TeV.

Note that photons emitted with energies above about 30 GeV can interact with stellar photons in the binary system to produce pairs $\gamma\gamma \rightarrow e^+e^-$ (Gould & Schröder, 1967). This leads at least to a hardening (softening) of the observed VHE γ -ray spectrum above (below) a few 100 GeV (Dubus, 2006). This can produce modifications to the electron distribution since pairs are created (this is further discussed in §6.1.1).

4.2. Luminosity

Assuming the situation is steady-state and that the pulsar spin-down power is largely given to the particles ($\sigma \ll 1$), the synchrotron and inverse Compton luminosities should be, to order unity, equal to \dot{E} . With a γ^{-2} injection, the inverse Compton and synchrotron luminosities are roughly equal (Moderski et al., 2005).

With such short radiative timescales, only a moderate spin-down power of $10^{36} \text{ erg s}^{-1}$ is sufficient to achieve a high radiative efficiency in X-rays and γ -rays. This has little to do with the assumptions made on the pulsar wind γ_p , σ or \dot{E} (if the wind is purely composed of electrons and positrons: a pulsar wind with a significant ion fraction would require a higher \dot{E} to obtain the same luminosity). The efficiency comes from

the high radiation and magnetic energy densities at the stand-off point which allow fast cooling of all of the injected particles. This contrasts with pulsars interacting with their supernova remnant or the interstellar medium, whose termination shock occurs much farther out, at correspondingly lower energy densities.

The main factor setting the high energy luminosity is the shock distance R_s or, alternatively, the orbital separation d_s (since $R_s \propto d_s$ in Eq. 5). The shock distance enters as R_s^2 in the synchrotron timescale compared to R_s in the escape timescale. The smaller R_s is, the more synchrotron emission dominates and the more the pulsar can be expected to be radiatively efficient. However, this is at the expense of the inverse Compton flux since the associated timescale stays constant with R_s .

The dominant radiation process does not change along the orbit for a given electron energy (because $\gamma_s \sim R_s/d_s$), but the probability that this electron can escape from the region of high U_\star and U_B without significant radiation increases with d_s (or R_s). The steady-state assumption then breaks down. Conditions change as particles flow away from the shock region. The overall nebula must then be taken into account to model the emission (§5).

4.3. Comparison to SEDs

The characteristic frequencies for all three γ -ray binaries were calculated using the parameters in Table 1 and additional assumptions on their stellar winds. For LS 5039, the wind is radiative with $\dot{M}_w = 10^{-7} M_\odot \text{ yr}^{-1}$ and $v_\infty = 2400 \text{ km s}^{-1}$. For LSI +61°303 and PSR B1259-63 two winds were assumed: a dense, equatorial outflow and a fast, polar wind similar to that in LS 5039. Details are given in §6. The standoff distances R_s and frequencies are given in Table 2.

Anticipating on §6, the pulsar was assumed to have $\gamma_p = 10^5$, $\sigma = 10^{-2}$ and $\dot{E} = 10^{36} \text{ erg s}^{-1}$ in all three systems. Note that \dot{E} and σ always appear in the same proportion so that for given R_s there are really only two parameters, γ_p and $\sigma\dot{E}$. The characteristic frequencies are not very sensitive to $\sigma\dot{E}$. Interestingly, the SED is also indifferent to the exact value of γ_p but for the low energy end of the inverse Compton spectrum.

The expected breaks match reasonably well the observed SEDs of LS 5039, LSI +61°303 and PSR B1259-63, notably the rising then flattening X-ray to γ -ray spectra. An important observable feature is that, all other parameters being equal, the break frequency in X-rays $\sim 1/d_s$ should be lower at apastron than at periastron. At higher energies, the break $\nu_{\text{comp},s}$ in LSI +61°303 and PSR B1259-63 occurs at frequencies a factor 2 lower than in LS 5039, because of the lower T_\star (Table 1). The TeV spectrum of PSR B1259-63 is indeed steeper than that of LS 5039. Unlike what happens with synchrotron, the spectral break in the inverse Compton emission should not change along the orbit if R_s is proportional to d_s (Eq. 5).

However, there are two shortcomings: the radio emission and the EGRET detections of LS 5039 and LSI +61°303 at GeV energies (Fig. 1). The radio emission cannot be addressed by the present scheme: the minimum synchrotron frequency is $\approx 1 \text{ keV}$ and radiation well below must originate from particles

that have cooled and escaped from the vicinity of the pulsar (see §5-6).

The GeV emission may be problematic as the model predicts a gap between 250 MeV and 50 GeV when $\gamma_p = 10^6$. There is no room for increasing the synchrotron emission so as to include at least a fraction of the EGRET band: the maximum synchrotron frequency is fixed by the assumptions on γ_{max} . Advocating higher γ_{max} seems unreasonable. Moreover, for large orbital separations as in PSR B1259-63 or LSI +61°303 (at apastron), the maximum electron energy is limited by the Larmor radius (supposed $\lesssim R_s$) rather than by radiative losses. These range from 2 MeV (apastron, polar wind) to 80 MeV (periastron, equatorial wind) for PSR B1259-63 and from 30 MeV (apastron, polar wind) to 250 MeV (periastron, equatorial wind) for LSI +61°303. The easiest solution consists in lowering the Lorentz factor γ_p of the particles in the pulsar wind to 10^5 . This lowers $\nu_{\text{comp},\text{min}}$ to a few GeV whilst leaving the spectral breaks unchanged. The inverse Compton timescale becomes long compared to t_{esc} at the low energy end of the electron distribution and, as for radio, high energy emission between 250 MeV and a GeV will also come from escaping particles.

5. A numerical model of the pulsar nebula

The detailed spectral energy distribution, including evolution of the physical conditions in the flow, is calculated here with the aim of accounting for the resolved radio emission as well as providing more quantitative comparisons. The assumptions are described in this section, with LS 5039 used as an illustrative example when necessary. The application and comparison with observations are made in §6.

5.1. Evolution of the particle distribution

Particles are continuously injected with a γ^{-2} power-law energy spectrum between γ_{min} and γ_{max} beyond the termination shock, in an emitting region of characteristic size R_s . Other distributions have not been considered for the sake of simplicity. The downstream particle density, in the limit of small σ is given by (Kennel & Coroniti, 1984b)

$$n_2 \approx 3n_1\gamma_p \approx 980 \gamma_6^{-1} \dot{E}_{36} R_{11}^{-2} \text{ particles cm}^{-3} \quad (24)$$

using Eq. 2. The normalisation of the injected particle power-law $Q(\gamma)$ is then found from

$$\int_{\gamma_{\text{min}}}^{\gamma_{\text{max}}} Q(\gamma) d\gamma \approx \frac{c}{3} n_2 \pi R_s^2 \approx 3 \cdot 10^{35} \gamma_6^{-1} \dot{E}_{36} \text{ particles s}^{-1} \quad (25)$$

Particles move downstream away from the pulsar with a speed v (initially $\approx c/3$) whilst emitting radiation. The resulting ‘cometary’ nebula is modelled by dividing it into small (logarithmic) subsections where the physical parameters (magnetic field, density etc) are homogeneous. The nebula is assumed to expand conically. The evolution of the distribution function $N(\gamma, z)$ along the nebula is given by

$$\frac{\partial N}{\partial z} = \frac{\partial}{\partial \gamma} \left[\frac{d\gamma}{dz} N \right] + Q(\gamma) \delta(z-1) \quad (26)$$

Table 2. Characteristic frequencies of the expected spectral energy distributions. The standoff distance (first number) is calculated at periastron and apastron using Eq. 3 and the stellar wind parameters in §6. Both a polar and an equatorial wind are considered for the Be stars. The break frequencies in the synchrotron and Compton spectra are then calculated using Eq. 19 and Eq. 22 in §4.1. The pulsar wind parameters are the same for all the objects: $\dot{E} = 10^{36}$ erg s $^{-1}$, $\sigma = 0.01$ and $\gamma_p = 10^5$. The star temperatures are such that inverse Compton interactions are in the Klein-Nishina regime.

LS 5039	periastron $d_s = 0.1$ AU			apastron $d_s = 0.2$ AU		
	$2 \cdot 10^{11}$ cm	100 keV	0.4 TeV	$4 \cdot 10^{11}$ cm	200 keV	0.4 TeV
LSI +61°303	periastron $d_s = 0.1$ AU			apastron $d_s = 0.7$ AU		
	polar	$8 \cdot 10^{11}$ cm	100 keV	0.8 TeV	$4 \cdot 10^{12}$ cm	15 keV
	equatorial	$5 \cdot 10^{10}$ cm	6 keV	0.05 TeV	$2 \cdot 10^{12}$ cm	5 keV
PSR B1259-63	periastron $d_s = 0.7$ AU			apastron $d_s = 10$ AU		
	polar	$3 \cdot 10^{12}$ cm	12 keV	0.5 TeV	$4 \cdot 10^{13}$ cm	1 keV
	equatorial	10^{12} cm	4 keV	0.2 TeV	not applicable	

where z is the distance along the nebula normalised to the standoff distance R_s . The radiative and adiabatic losses for an electron (positron) are:

$$\frac{d\gamma}{dz} = \frac{R_s}{v} \left[\left(\frac{d\gamma}{dt} \right)_{\text{sync}} + \left(\frac{d\gamma}{dt} \right)_{\text{comp}} + \left(\frac{d\gamma}{dt} \right)_{\text{ad}} \right] \quad (27)$$

The synchrotron radiation loss is the usual $\dot{\gamma}_{\text{sync}} \sim \gamma^2 B^2$. The energy loss to inverse Compton radiation is calculated using the Jones (1968) kernel, which is a continuous approximation although electrons can lose significant amounts of approximation in discrete encounters with photons in the Klein-Nishina regime. Furthermore, the Jones kernel supposes an isotropic photon distribution. This should not be too bad an approximation as the angle-dependence of the cross-section is limited in the Klein-Nishina regime.

The adiabatic losses for two dimensional expansion of relativistic material are

$$\left(\frac{d\gamma}{dt} \right)_{\text{ad}} = \gamma \left(\frac{d \ln n^{1/3}}{dz} \right) = -\frac{2}{3} \frac{\gamma}{z} \frac{v}{R_s} \quad (28)$$

The evolution of the physical parameters of the flow (B , v etc) with z is described in the next section. Once these are determined, the differential equation governing the evolution (Eq. 26) is numerically integrated along z using a logarithmic grid in γ , following the scheme proposed by Chang & Cooper (1970). The results presented thereafter typically use at least 1000 points for the particle energy grid and 200 points in frequency ν to represent emission spectra. The results have been checked to be independent of the numerical details.

5.2. Evolution of physical conditions in the flow

The evolution of the nebular flow speed v and magnetic field B with z are computed by using the relativistic Bernoulli equation stating that

$$\frac{\Gamma}{n} \left(\frac{B^2}{4\pi\Gamma^2} + P + ne \right) = \text{constant} \quad (29)$$

along streamlines for stationary, adiabatic motion. Γ is the bulk Lorentz factor of the flow $\Gamma = (1 - v^2/c^2)^{-1/2}$. The additional relationships for the magnetic field B , density n , pressure P and internal energy e come from the conservation of (toroidal) magnetic flux Bvz/Γ , of the number of particles nvz^2 and the relativistic equation of state $P = n(e - mc^2)/3 \propto n^{4/3}$ (Kennel & Coroniti, 1984a).

Close to the shock, the total energy is dominated by pressure. The density is then roughly constant, the flow speed decreases as $v \propto 1/z$ and the magnetic field increases because of flux conservation (Fig. 3). The flow becomes magnetically dominated at large radii ($z \gtrsim 10$) at which point the density varies as $1/z^2$, the magnetic field decreases as $B \propto 1/z$ and the asymptotic flow speed is σc (Kennel & Coroniti, 1984a).

The present approach is admittedly simplistic. MHD simulations of a spherically symmetric pulsar wind interacting with a constant external medium exist (Bucciantini et al., 2005) and give insights into possible shortcomings. The shape of the pulsar termination shock is bullet-shaped rather than spherical. An asymmetric pulsar wind would surely change the aspect of the flow (as it does with the Crab and Vela pulsars). Shocked material flows along two channels, a fast one with material from the forward termination shock and a slower one with material from the backward facing termination shock. The nebular flow loses energy to radiation, contradicting the adiabatic approximation. Collimation shocks may re-energise particles along the jet; mixing can occur at the surface discontinuity between the fast-moving shocked pulsar wind and the slow-moving stellar wind. Mass loading will slow down if not disrupt the flow. At larger distances, the flow shears because of orbital motion (§5.4), leading to additional complications. Finally, the flow merges with the stellar wind and/or ISM on scales of ~ 0.01 pc based on the densities.

5.3. Calculated spectrum

Neglecting any effect of the orbital motion (see §5.4), the overall spectrum results from the summed contribution of the syn-




Fig. 3. From left to right: evolution in the pulsar nebula of the flow speed, magnetic field and density as a function of the normalised distance z . The rightmost panel shows the time needed to reach a given distance z in the flow. At $z \approx 800$ ($d = zR_s \approx 10$ AU), the elapsed time is equal to the orbital period (dotted line). The model parameters are $\dot{E} = 10^{36}$ erg s $^{-1}$, $\gamma = 10^5$, $\sigma = 0.01$ and $R_s = 2 \cdot 10^{11}$ cm, appropriate for LS 5039 at periastron.

chrotron and inverse Compton spectra (on stellar photons) calculated for subsections dz of the nebula extending away from the star. As was done for energy losses, the particle distribution and radiation fields are assumed spatially homogeneous and isotropic in each subsection. The inverse Compton spectrum takes into account the reduction of the cross-section for interactions in the Klein-Nishina regime via the Jones kernel. The radiation field of the stellar companion is simply diluted by the inverse square of the distance to the star ($U_\star \propto d_\star^{-2}$ with $d_\star = d_s + (z - 1)R_s$).

Fig. 4 illustrates the evolution of the emission spectrum and particle distribution with increasing distance z to the pulsar in the case of LS 5039. The first 3-4 spectra show the initial evolution towards the steady-state described in §4. The particle distribution above $\gamma_s \sim 10^6$ cools rapidly at high energies due to synchrotron losses, resulting in a plateau down to ν_s as expected. Evolution then sets in around $z \sim 10$, with the spectral luminosity decreasing with B . The particle distribution steepens with emission sweeping the optical to IR bands. Further evolution, corresponding to peak emission in mm to radio, is set by adiabatic losses.

Self-Compton radiation is negligible. Synchrotron emission was optically thin down to low radio frequencies in all the models computed for §6. This is a consequence of the flow geometries, which are discussed in §5.4. Still, synchrotron self-absorption might be possible in a flow directed predominantly along the line-of-sight.

In the absence of orbital motion, the spectrum seen by the observer is simply the integrated emission along the ‘comet’ tail nebula, just as for isolated pulsar interacting with the ISM. However, as can be deduced from Fig. 4, the radio emission originates from regions several AU away from the pulsar ($z \gtrsim 100$). The time Δt it takes for particles to reach this location, about a day here, is a significant fraction of $P_{\text{orb}} = 3.9$ days. Orbital motion will have changed the inner boundary condition so the spectrum seen by an observer will be a composite of SEDs from all orbital phases ϕ . Here, the only parameter dependent on the orbital phase is the pulsar standoff point R_s . Other dependences can be envisioned if, for instance, the pulsar wind is anisotropic or if the magnetization parameter is a func-

tion of distance to the light cylinder (with higher σ expected close to the pulsar).

The relevant timescales to compare are the flow timescale $\tau_{\text{flow}} = d/v$ to reach a distance d from the shock and the orbital motion timescale $\tau_{\text{orb}} = d_s/v_{\text{orb}}$. The orbital timescale can vary greatly for a highly eccentric orbit. Nevertheless, taking v equal to its asymptotic value σc , the distance where orbital motion becomes important will on average be

$$d_n \approx \sigma c P_{\text{orb}} / 2\pi \quad (30)$$

and the associated timescale, $P_{\text{orb}}/2\pi$. As it turns out, the synchrotron emission from LS 5039 depends little on the orbital phase because R_s/d_s is constant with ϕ . The spectrum seen by an observer should therefore be close to the integrated spectrum shown in Fig. 4 even at low frequencies (see §6.1).

5.4. Appearance of the cometary nebula

On a small scale, the shocked material flows away from the binary on a straight path, following the direction given by the vector difference of the stellar wind and orbital speeds $\mathbf{v}_w - \mathbf{v}_{\text{orb}}$. On a larger scale, the material ‘bends’ as the orbital motion becomes important at $\sim d_n$ (about 1 AU for LS 5039; Eq. 30). Parcels of material are still on outward straight paths but the orbital motion shears the nebula. The opening angle of the flow ($\sim R_s/d_s$) is small so that the width of the ‘comet’ tail is smaller than d_n . For a circular orbit, the large scale appearance is then an Archimedes spiral with a step size $2\pi d_n = \sigma c P_{\text{orb}}$.

More generally, the flow speed amplitude and direction change along the orbit and the exact shape can differ from a perfect spiral. Furthermore, emission along the spiral is not constant but depends upon the distance d to the pulsar or, alternatively, upon the time elapsed since the particles left the pulsar. The flux received by the observer may also be Doppler by factors of a few (de)boosted initially, when the flow speeds are high (neglected here). Finally, the orbital orientation will also change the appearance of the nebula. For an edge-on system ($i = 90^\circ$), the nebula will mimic a microquasar bipolar jet.

The trajectory of material leaving the pulsar wind termination shock can be found as a function of the elapsed time t ,




Fig. 4. Left: Evolution of the emission along the pulsar nebula. The pulsar nebula is divided into sections at various distances z with the corresponding synchrotron (black) and inverse Compton (grey) spectra shown. The first section shown is for $z = 1$ ($dz = 0.01$). The section dz is then increased: the fifth spectrum is at $z \approx 1.5$, the tenth at $z \approx 15$, the last spectrum shown is at $z \approx 1000$. The total spectrum of the pulsar nebula is shown as a thick black line. The model parameters are the same as in Fig. 3. The thermal emission at ~ 1 eV is direct light from the stellar companion. Right: Evolution of the particle distribution as the shocked pulsar wind moves out from the system. The sections are the same as those shown in the SED. The initial γ^{-2} power law cools rapidly at high energies due to synchrotron radiation. Adiabatic losses dominate at large distances.

using polar coordinates (r, θ) in the orbital plane. The distance d reached along the nebula is

$$d = \int_0^t v dt \quad (31)$$

where v is the flow speed calculated in §5.2. The radius r with the coordinates centered on the massive star is given by

$$r^2 = d^2 + 2dd_s \cos \beta + d_s^2 \quad (32)$$

where β is the angle at which the shocked material flows and d_s is the star-pulsar separation at $t = 0$. If \mathbf{e}_v is the unit vector along the flow direction and \mathbf{e}_r is the radial unit vector from the massive star to the pulsar location at $t = 0$, then $\cos \beta = \mathbf{e}_r \cdot \mathbf{e}_v$. The angle θ of the material is then

$$\theta = \omega + \eta - \psi \quad (33)$$

where ω is the periastron argument, η is the true anomaly of the pulsar at $t = 0$ (angle measured from the star center, $\eta = 0$ at periastron) and ψ is derived from $r \sin \psi = d \sin \beta$.

The full pattern follows from having η vary with time t according to the usual relationships governing orbital motion (*i.e.* using the eccentric anomaly). Note that v , d_s and β also vary with orbital phase. The result is then projected onto the plane of the sky, given the orbital inclination i .

Fig. 5 shows the spiral pattern computed for LS 5039 with $\dot{E} = 10^{36}$ erg s $^{-1}$, $\gamma = 10^5$, $\sigma = 0.01$ and the pulsar located at periastron. The figure represents the emission from particles ‘ejected’ over a timescale of about an orbital period and a half. The spiral ‘turns’ around and outwards as the pulsar moves along the orbit. The lines illustrate the direction at which material flows out from the binary. Spacing between lines correspond to equal intervals of time; in other words, lines are more spaced out at periastron due to fast orbital motion (before projection on the plane of the sky). For very eccentric orbits (such

as PSR B1259-63, see §6.3), material will preferentially flow out in the direction of the major axis on the apastron side (for a radial outflow). There will be a preferred emission axis, even when the system is seen pole-on. Another effect illustrated by Fig. 5 is the bunching of lines due to the projection on the sky plane. A binary with a circular orbit but seen at some inclination will also have a preferred emission axis.

Besides the orbital eccentricity and inclination, a third effect that will shape the appearance of the cometary nebula is the variation of emission with orbital phase. The amplitude can change with orbital phase. But even if the spectrum varies little along the orbit, the locus of peak emission changes as the elapsed time depends directly on the absolute scale R_s (hence phase). If the standoff point is twice as far away at apastron than it is at periastron, then the spatial scale on which the radio peaks also doubles. In Fig. 5, the intensity of the 5 GHz radio emission is represented by the grayscale coloring. Note how far out material on the upper left hand side (corresponding to apastron) has a higher intensity than material on the right hand side (periastron) although the latter was emitted a shorter time ago.

All these effects combine to create radio maps that are far from trivial (see §6). Despite the complexity, there is one simple prediction: the step of the spiral should be $\approx \sigma c P_{\text{orb}}$. In principle, the radio nebula of γ -ray pulsar binaries could be used to measure the average magnetization parameter of relativistic pulsar winds. The particular cases of LS 5039, LSI +61°303 and PSR B1259-63 are examined in the next section.

6. Application to the observed γ -ray binaries

6.1. LS 5039

The wind of the O6.5V companion star in LS 5039 is assumed to be radiatively-driven with a velocity and density given by

$$v_w = v_\infty \left(1 - \frac{R_\star}{d_s}\right)^\beta \quad \rho_w = \frac{\dot{M}_w}{4\pi d_s^2 v_w} \quad (34)$$

with $\beta \approx 1$. Spectroscopic observations of LS 5039 constrain \dot{M}_w to $\approx 10^{-7} M_\odot \text{ yr}^{-1}$ and $v_\infty \approx 2400 \text{ km s}^{-1}$ (McSwain et al., 2004). The wind is assumed isotropic and purely radial. Since the orbit of the pulsar takes it to almost a stellar radius of the companion at periastron in LS 5039 and LSI +61°303, more complex wind launching conditions than those assumed above could play a role in setting R_s . A significant wind rotation might also influence the location of the standoff point and the flow direction.

The pulsar spindown power was taken to be $\dot{E} = 10^{36} \text{ erg s}^{-1}$, close to that measured in PSR B1259-63 (§6.3). R_s then varies between $2 \cdot 10^{11} \text{ cm}$ (periastron) and $4 \cdot 10^{11} \text{ cm}$ (apastron). The two parameters left are the pulsar wind magnetization σ and Lorentz factor γ_p . Low values of $\sigma \gtrsim 10^{-4}$ yield low downstream magnetic fields. Hence, the inverse Compton emission dominates with far too little X-rays emitted to match the SED. At the other end, values of $\sigma \approx 0.1 - 1$ lead to overall emission inefficiency because electrons move quickly out into regions of low radiation fields (the downstream flow speed is high). Values of σ in the 0.001-0.01 range yield the best results. However, with $\sigma \approx 0.001$, the characteristic size of the radio emission (Eq. 30) would be too small by a factor 10 to match the observed scale of $\sim 10 \text{ AU}$ (Paredes et al., 2000, see below). The best compromise was $\sigma = 0.01$. Coincidentally, the downstream flow speed saturates at a value ($\sim 3000 \text{ km s}^{-1}$) that is then close to the terminal velocity of the stellar wind ($\sim 2000 \text{ km s}^{-1}$).

The Lorentz factor γ_p influences the lower cutoff of the particle distribution. Lower values enable the inverse Compton emission to extend down to the GeV range, as discussed in §4.3. With more low energy electrons, the radio emission is also more easily accounted for. A value of $\gamma_p = 10^5$ was the best compromise: at 10^6 the GeV emission is unaccounted for, at 10^4 the radio emission is overpredicted and the downstream particle distribution then extends more than four decades in energy. The adopted values are $\sigma = 0.01$ and $\gamma_p = 10^5$.

6.1.1. The SED

The evolution of the SED as the material flows away from the pulsar is shown in Fig. 4. The high energy SED is very much that described in §4, with the spectral breaks in the synchrotron and inverse Compton spectra at their expected locations. Electrons in the comet tail enable the summed emission to extend down to low frequencies and also contribute to emission in the GeV domain. The comparison to the observations shows reasonable but not perfect agreement. Changes in the particle distribution, stellar or pulsar wind could reduce the discrepancies but have not been attempted here. The purpose

at this stage is to validate the model without fine-tuning and to compare the γ -ray binaries under similar assumptions.

Fig. 6 shows the variation of the SED between periastron and apastron. Apart from the slight lowering of the synchrotron break frequency (see §4.3), the spectral energy distributions are nearly identical. This agrees with the lack of strong orbital variability observed in X-rays or radio. However, an orbital modulation is unavoidable at TeV energies because of $\gamma\gamma$ absorption (see below).

At γ -ray energies, the model has plateau emission across the EGRET band and up to the HESS regime, as expected in §4. However, (1) the GeV flux is underestimated and (2) the high energy break is a factor 2-3 too low than what would be required to match the hard HESS power-law ($\sim 2 \text{ TeV}$). LS 5039 is in the Galactic Plane so that the EGRET flux could be overestimated (i.e. should be interpreted as an upper limit) if the diffuse γ -ray background subtracted is underestimated or if several unresolved sources contribute to the emission. That the measurements are non-contemporaneous should not be a problem in the present scenario, where variability is expected to be small (except for the possible pair cascade, see below).

The approximate treatment of the emission geometry may be responsible for these discrepancies. Adjusting \dot{E} , σ or the standoff point distance can alleviate both problems (Eq. 22). Tweaking the accelerated distribution of particles might help, but lacks justification. There could be a direct contribution from acceleration processes in the pulsar magnetosphere. Two other avenues can also be explored.

First, the anisotropy of the stellar radiation field might play a role. The outgoing energy of an up-scattered photon depends on the interaction angle with the higher energies reached at π . Even if the distribution of electrons is isotropic, the mean angle of interaction of the photons detected by the observer will depend on the orbital phase, hence there should be an energy effect. This is likely to be small at TeV energies as there is not much energy dependence on angle in the Klein-Nishina regime (Moderski et al., 2005).

A second important effect is $\gamma\gamma$ absorption (Böttcher & Dermer, 2005; Dubus, 2006; Bednarek, 2006). High energy γ -rays are emitted in the immediate vicinity of the pulsar where they can then interact with stellar photons to produce e^+e^- pairs. The threshold γ -ray energy is $\approx 30 \text{ GeV}$. The transmitted flux is shown in Fig. 6 as grey lines for both periastron and apastron. The calculation assumes that γ -ray emission is isotropic and occurs at the pulsar location (see Dubus 2006 for details). The opacity depends on orbital phase, with a maximum close to periastron passage when most of the γ -ray emission between 0.1-10 TeV should be absorbed. An orbital modulation of the VHE flux measured by HESS is therefore expected. Note that this is not a discriminating feature with the accretion/jet scenario since any γ -ray emission occurring within $\approx 1 \text{ AU}$ of the companion star in this system (for instance, in regions close to the black hole) would be significantly modulated.

The average result of absorption will be a hardening of the high energy (HESS range) spectrum. Pairs created by the absorption of γ -rays can subsequently upscatter stellar photons to very high energies, thereby initiating a cascade (not taken

into account in the calculation shown in Fig. 6). This can significantly increase the emission below 30 GeV as the absorbed energy is redistributed below the interaction threshold if the cascade is very strong (Aharonian et al., 2006). This would also introduce orbital variability below 30 GeV (Dubus, 2006; Bednarek, 2006). However, note that high energy pairs above γ_s will radiate predominantly synchrotron emission instead of upscattering photons (§3.4), reducing the extent of the cascade (Aharonian et al., 2006).

Understanding the exact impact of these two effects will require complex modelling, including a better representation of the shock geometry. The conclusion is that the present fairly rustic model is already able to account for the whole SED adequately, in the manner described in §4 using straightforward arguments.

6.1.2. Radio maps

The total radio intensity does not vary much along the orbit but the time/distance at which maximum radio emission is reached does depend on the orbital phase. The evolution of the radio flux with distance to the pulsar was calculated as above for various orbital phases and used together with the geometric model for the nebula (§5.4) to create emission maps of the system. Free-free absorption on the stellar wind becomes optically thin at ~ 1 AU (see rough estimate in §2.4), close to the distance at which radio emission starts. Hence, this effect is neglected.

Fig. 7 shows the evolution of the 5 GHz emission as a function of orbital phase ϕ . The first map (periastron $\phi = 0$) shown uses the same parameters as those in Fig. 5, except that the emission is smoothed by a Gaussian with FWHM of 0.8 mas to approximate the observing setup of Paredes et al. (2000). The system is seen at an inclination $i = 120^\circ$, i.e. 30° below the orbital plane ($i = 60^\circ$ would be 30° above the plane, the degeneracy from the radial velocity is lifted by the comparison of the curvature in the map to the VLBI observations). The contours at 50%, 25% and 10% of the maximum surface intensity can be compared to the 5 GHz VLBI observation. The maps at $\phi = 0 - 0.25$ are strikingly similar to that observed by Paredes et al. (2000), with slightly curved emission extending a few mas and the inner emission pointing in a slightly different direction. The observed very weak counter-jet would be associated with emission emitted at the previous periastron passage.

The model predicts changes in the radio maps with ϕ (Fig. 7) that can be verified by further VLBI observations. An animated map gives the distinct impression that blobs are alternatively ejected to the sides of the system, cunningly mimicking a microquasar. Accurate astrometry should show that the position of peak radio emission varies by a few mas. The orbital motion produces a ‘comma’ shape that is more pronounced at $\phi \approx 0$, when particles emitted at *apastron* start emitting in the radio band. Emission occurs principally along the major axis of the sky-projected system orbit. This effect shows up as a bunching of the lines representing the trajectories in the mid-plane of Fig. 5.

On progressively larger scales the emission becomes egg-shaped along the same axis, with the tip of the ‘egg’ on the

apastron side as projected onto the sky. MERLIN and EVN radio observations do resolve emission on a larger scale on one side than on the other (Paredes et al., 2002). The observed emission is more linear than in the model with $i = 60^\circ$. A greater inclination would help since emission becomes exactly line-shaped for an edge-on orbit $i = 90^\circ$. However, the X-rays would then be eclipsed when the pulsar moves behind the companion; an upper limit using the orbit and assuming the X-ray emitting region is a sphere of radius R_s gives $\approx 75^\circ$. The observed extent of radio emission is also larger than expected. On an equivalent map to the EVN observation, the 5 GHz intensity decreases to 1% of the peak value at 12 mas on one side and 6 mas on the other instead of 35 mas and 25 mas. The quantitative agreement is not as good as on scales $\sim \sigma c P_{\text{orb}}$.

While the map in Fig. 7 shows basically only emission within an orbital timescale, maps on scales $\gg \sigma c P_{\text{orb}}$ sum up emission from dozens of orbits, representing only a tiny fraction of the initial energy budget. The physics on such scales is undoubtedly more complex than the simple approach taken in §5.2: mixing occurs with the surrounding medium; recollimation shocks re-energise the electrons; shearing introduces turbulence; magnetic field changes as it is azimuthally stretched; shock conditions at earlier times were different etc. Note that the outflow length of ~ 500 AU in Paredes et al. (2002) coincides roughly with the distance at which the stellar wind merges with the ISM. Incidentally, many of these complications are also encountered in the propagation of relativistic jets.

The conclusion is that the pulsar model provides a very good quantitative explanation for the radio emission emitted on scales of AU in LS 5039 and provides a qualitative framework to understand radio emission on larger scales. Radio observations have tremendous potential to validate the model.

6.2. LSI +61°303

The pulsar \dot{E} and γ_p were taken similar to those of LS 5039 for comparison purposes. A slightly higher value of $\sigma = 0.02$ was adopted here to better match the observed radio scales (§6.2.4; note that Table 2 uses $\sigma = 0.01$).

6.2.1. Stellar Wind

The stellar companion in LSI +61°303 is a B0Ve type star which significantly complicates modelling over the case of LS 5039. Ultraviolet and infrared observations of Be stars show that their stellar wind has two components: a fast, polar outflow with properties typical of radiatively-driven winds; and a slow, equatorial outflow (Waters, 1986).

A typical polar wind was modelled using Eq. 34 with $\dot{M}_w = 10^{-8} M_\odot \text{ yr}^{-1}$ and $v_\infty = 2000 \text{ km s}^{-1}$ (Waters et al., 1988). The standoff point R_s then varies between $8 \cdot 10^{11} \text{ cm}$ and $4 \cdot 10^{12} \text{ cm}$. The equatorial wind was modelled as

$$\nu_w = \nu_0 (R/R_\star)^{n-2} \quad \rho_w = \rho_0 (R/R_\star)^{-n} \quad (35)$$

with $n \approx 3$, $\nu_0 \approx 10 \text{ km s}^{-1}$ and $\rho_0 \approx 10^{-11} \text{ g cm}^{-3}$ from the comparison to infrared observations of LSI +61°303 (Waters et al., 1988; Martí & Paredes, 1995). This corresponds




Fig. 7. Orbital evolution of the 5 GHz radio emission from the pulsar nebula in LS 5039. The maps were composed by summing up the contributions along the spiral nebula, as calculated for the various orbital phases ϕ (see Fig. 5 for the $\phi=0$ case). The maps were then smoothed by a gaussian kernel with a Full-Width at Half Maximum (FWHM) of 0.8 mas ($3 \cdot 10^{13}$ cm at 2.5 kpc). The maps share a common color scale to allow direct comparison, with contours at 50%, 25% and 10% of the maximum orbital radio emission. The maps at $\phi=0-0.25$ are very similar to that observed by Paredes et al. (2000).

to an equatorial mass flux about a 100 times larger than the polar mass flux. The typical opening angle of the Be star equatorial outflow is 15° . $H\alpha$ observations show the disc extends to $5-20 R_\star$ and that the terminal velocity is a few hundred km s^{-1} . The standoff point is then between $5 \cdot 10^{10}$ cm and $2 \cdot 10^{12}$ cm, smaller than the polar wind values because of the higher densities. The major difference between the two types of stellar winds will be at periastron.

6.2.2. The SED

The resulting SEDs at periastron and apastron for both stellar winds are shown in Fig. 8. They agree well with the expectations set out in Table 2, most notably the much lower break frequencies for the equatorial outflow compared to the polar wind due to the smaller R_s .

Orbital variations in X-rays are expected to be greater than in LS 5039. Indeed, observations show variations of a factor ~ 10 in soft X-rays (0.1-2 keV) and of smaller amplitude at higher energies. The maximum is inferred to be close to periastron (Harrison et al., 2000; Casares et al., 2005a). Consulting Fig. 8, this would seem to favor combining a small R_s at periastron (equatorial wind) and a large R_s at apastron (polar wind). At apastron, the pulsar is $15 R_\star$ away from its stellar companion, in a region where the Be equatorial disc could already be truncated.

At γ -ray energies, the TeV fluxes are consistent with the Whipple upper limits. Small R_s for the equatorial flow result in high magnetic fields, conversely lowering the γ -ray emission. Non-intuitively, TeV emission will be higher at *apastron* than at *periastron* if the pulsar plows through a dense equatorial outflow close to its companion and a more tenuous wind further out. With R_s closer to the pulsar, synchrotron losses largely dominate at periastron over IC emission. This can be tested by observations with VERITAS or MAGIC. As with LS 5039, the GeV fluxes are too low to match the EGRET spectrum. An e^+e^- cascade is unlikely to play a role here as the $\gamma\gamma$ opacity is small except very close to periastron (Dubus, 2006).

At low energies, the radio emission is too steep, primarily because of the value of σ . A lower value ($\sigma \approx 0.001$) would

yield better agreement: the downstream magnetic field is then lowered, reducing synchrotron losses and enabling stronger radio emission at large z . However, the physical extent of the associated radio emission would be smaller. A compromise would be to have σ change with orbital phase. The distance of the shock to pulsar R_s changes by an order-of-magnitude here (compared to a factor 2 in LS 5039) and σ could be envisioned to decrease as R_s increases.

6.2.3. Radio outbursts

LSI +61°303 has regular radio outbursts during which the flux increases by an order of magnitude. The outbursts last several days, peaks at an orbital phase $\approx 0.2 - 0.7$ and repeat on the orbital period. There is also an X-ray orbital modulation (a factor 10 in the *ROSAT* band, less above) that precedes the radio peak by $\Delta\phi \approx 0.1 - 0.5$ (the exact delay varies between outbursts). PSR B1259-63 has similar outbursts at periastron, associated with the pulsar crossing the Be equatorial outflow (Melatos et al., 1995). Indeed, the Be nature of the companion is the main feature distinguishing PSR B1259-63 and LSI +61°303 from LS 5039, which has no radio outbursts.

Assume the pulsar samples the dense equatorial disc around periastron, and a more tenuous, fast wind at apastron. Changes in R_s , possibly σ , are expected along the orbit; yet these do not suffice to explain the outbursts because of the timescales involved. If such was the case, the ~ 10 -day delay in the phasing of the X-ray and radio modulations would presumably be associated to the time it takes for electrons to cool from injection. The strongest radio emission would occur at apastron, when the shock is far from the pulsar and σ is expected to be lower. The elapsed time to radio peak is then greater than the orbital period (grey line in Fig. 8) and this would smooth out the flux modulation.

The impact of the Be star equatorial outflow is probably somewhere else. With a slow wind, the standoff point $R_s \approx 5 \cdot 10^{10}$ cm is very small at periastron. Moreover, the equatorial wind is very slow so that the Bondi accretion radius $\propto 1/v_w^3$ (§2.4) is actually greater than R_s . For the adopted parameters, the accretion radius is $\approx 2 \cdot 10^{11}$ cm at periastron. Matter falls

towards the compact object at a rate $\approx 10^{17} \text{ g s}^{-1}$, but is not accreted: the pulsar wind is still able to hold off the inflow of material (§2.4). However, the situation changes in a significant way from that described until now.

When the relative motions are fast (polar wind case, or high velocity isolated pulsar case), the standoff distance is farther out than the Bondi capture radius so that the external ram pressure is mostly unidirectional. The pulsar creates a long comet tail directed away from the maximum ram pressure point. At periastron, the standoff distance is within the Bondi capture radius. The external pressure from inflowing material is more spherically distributed, as in a SNR plerion, so that the shocked pulsar wind cannot break open the surrounding medium on a short timescale.

Therefore, a possible picture would be that the shocked pulsar wind is embedded in the equatorial disc at periastron (plerion-like) and that the geometry is open (comet-like) at apastron, where the Bondi radius is smaller than R_s for both an equatorial or a polar wind. High energy radiation is not affected much as it is emitted close to the pulsar in both cases. The change should be mostly in the low frequency emission as particles are constantly injected in a bubble slowly expanding in the equatorial outflow. The outburst would be associated with emission from this bubble of electrons, following an earlier suggestion by Leahy (2004). Detailed modelling is not attempted here but note that adiabatic expansion of bubbles has been shown to reproduce the radio outbursts (Taylor & Gregory, 1984; Paredes et al., 1991; Leahy, 2004) and, naturally, has also been proposed for the radio outbursts of PSR B1259-63 (Connors et al., 2002).

6.2.4. Radio map

The radio emission map expected from the simple case of a polar wind is shown in Fig. 9 for $\phi = 0.5$ and $i = 120^\circ$. (Here, $\phi = 0$ corresponds to periastron passage, to which ≈ 0.2 should be added if using the radio ephemeris, Gregory 2002.) The orientation of the orbit is illustrated in Fig. 2 of Casares et al. (2005a). The image does not take into account the radio outbursts but should nevertheless be indicative of the overall morphology to expect.

As in LS 5039, the nebula has a ‘comma’ shape. The physical scale is larger than in LS 5039 because of the longer orbital period. The major axis is almost in the plane of the sky, defining a very clear preferred direction for emission. This is consistent with observations of one-sided radio emission on scales of 10-100 mas (Massi et al., 2001, 2004). The mean position angle will increasingly vary as one goes to higher spatial resolutions and the impact of the orbital motion becomes clearer. This could explain the almost perpendicular directions seen at 3 mas and 10 mas resolution, albeit at different epochs (Massi et al., 1993, 2001). Massi et al. (2004) also find changes in structure about 50 mas away from the core emission on timescale of days during the radio outburst. This could be due to a combination of rapidly changing emission angles at periastron passage and delayed peak radio emission; a proper model for the radio outbursts would be required to verify this possibility.

6.3. PSR B1259-63

For PSR B1259-63 the measured \dot{E} is $8 \cdot 10^{35} \text{ erg s}^{-1}$ (Manchester et al., 1995). The other pulsar parameters were taken to be $\sigma = 0.01$ and $\gamma_p = 10^5$ to compare with the other two systems. Emission by containment of a pulsar wind in PSR B1259-63 has previously been investigated by several authors (Tavani et al., 1994; Melatos et al., 1995; Tavani & Arons, 1997; Kirk et al., 1999; Murata et al., 2003). The purpose here is to put the system in the same context as LS 5039 and LSI +61°303, and explore possible large scale radio emission.

Both an equatorial and a polar stellar wind were considered, with the same parameters as those given for LSI +61°303 in §6.2.1. The periastron distance in PSR B1259-63 is comparable to the apastron distance in LSI +61°303. Any equatorial outflow is then at or close to its terminal velocity and there is little difference in R_s with a polar wind (only a factor 3, see Table 2). The equatorial wind is truncated at $\sim 15 - 20 R_\star$ and is not considered at apastron. Only a polar wind can contain the pulsar wind at apastron.

6.3.1. The SED

The spectral energy distribution at periastron and apastron is shown in Fig. 10 for the case of a polar wind. The SED at periastron for an equatorial wind (not shown) is very close to the one shown here, the difference in R_s being small. Unsurprisingly, the SED is essentially the same as that for LSI +61°303 at apastron, the orbital separation being the same and other parameters varying little. GeV emission from PSR B1259-63 around periastron should be detected by *GLAST* in the future, if only because of this similitude. The TeV slope appears in good agreement with the HESS measurement. The variations in integrated flux seen by HESS around periastron are beyond this model’s present abilities (which predicts none) and will probably require taking into account anisotropic IC scattering (see Kirk et al., 1999), as well as changes in the circumstellar density to properly reproduce. There seems to be an inflection in the X-ray spectrum between 1-10 keV that might correspond to the break observed with *RXTE*. Very good agreement can indeed be obtained when using $\gamma_p = 10^6$.

The SED at apastron is only slightly different. Despite the considerably larger orbital separation, the TeV flux is only lowered by factors of a few. The photon density is indeed much lower, but the scale of the flow (set by R_s) is much larger than at periastron. Particles have a longer time available to interact and this compensates for the weaker densities. Overall, the radiation efficiency is the same as at periastron as the same amount of energy is being dissipated regardless of orbital phase. The model cannot account in its present simplified form for the observed order-of-magnitude changes in X-ray luminosity. This was noted by Murata et al. (2003) who proposed that lowering σ with pulsar distance might help.

Emission at apastron occurs on large, resolvable scales. The grey dashed lines in Fig. 10 show the amount of the total flux emitted after an elapsed time $\Delta t > P_{\text{orb}}$ after injection. This flux

is radiated several hundreds of AUs away and may lead to a faint arcsecond scale bowshock structure up to IR frequencies.

6.3.2. Radio outbursts and map

In PSR B1259-63, observations of the radio dispersion measure imply the disc is inclined with respect to the orbital plane (Melatos et al., 1995). The radio outbursts seen before and after periastron are presumably associated with the two pulsar disc crossings (Ball et al., 1999). Working by analogy, smaller separations and a coplanar equatorial outflow would explain the single outburst per orbit in LSI +61°303. As explained above for LSI +61°303, the outbursts are unlikely to be due solely to a change in R_s . The large increase in radio flux might be related to the slow expansion of a bubble of relativistic electrons formed in the dense flow. Shock-acceleration of electrons from the stellar wind itself may also play a role (Ball et al., 1999).

It is therefore as difficult as with LSI +61°303 to accurately predict the shape the radio emission might take. Fig. 11 shows expectations at $\phi = 0.1$ assuming PSR B1259-63 only interacts with a polar wind. The spatial scales should be easy to resolve since R_s is large. Unfortunately there are no published radio maps to compare with. The rapid revolution of the pulsar around periastron produces an arc with two brighter ‘knots’ where pre- and post-periastron electrons are cooling. These two knots gradually drift apart. This takes no account of the radio outbursts. If these are localised in the equatorial wind, the impact might be only an increase in the central (unresolved) component, leaving the much fainter diffuse components described here mostly unchanged. The bow shock points away from the star in almost the same direction for most of the highly eccentric orbit.

7. Conclusion

The pulsar wind scenario offers a tantalizing way to unite LS 5039, LSI +61°303 and PSR B1259-63 into a new-class of young, rotation-powered γ -ray binaries. At present, nothing rules out that the compact object in LS 5039 and LSI +61°303 is a pulsar. The radial velocities allow it and there are no conclusive signs of accretion. Directly unveiling a radio pulse will not be possible due to strong free-free absorption; although a difficult endeavour, an X-ray or γ -ray pulse would prove the scenario.

A similar relativistic pulsar wind with most of its $\dot{E} \approx 10^{36}$ erg s⁻¹ energy ($\sigma \approx 0.01$) in electrons and positrons of Lorentz factor $\gamma_p \approx 10^5$ suffices to explain most properties in all three objects. The termination shock occurs at the pressure balance point. There, particles in the pulsar wind are scattered and accelerated into a non-thermal power-law distribution. The maximum energy is set by the size of the shock region and radiative losses. Downstream values in the shocked pulsar wind material (density, magnetic field) are set by the MHD jump conditions. Particles radiate synchrotron and inverse Compton whilst being advected away in a comet tail nebula. At high energies, synchrotron dominates over the Klein-Nishina cross-section, giving characteristic break frequencies between a rising and flat νF_ν spectrum in hard X-rays, and between a flat

and a declining νF_ν spectrum in VHE γ -rays. The VHE emission occurs close to the pulsar and an orbital modulation of the flux in the TeV range is expected in this scenario, due to absorption (pair production) on stellar photons.

The small-scale radio emission is also reasonably reproduced. Radiation shifts to lower wavelengths as particles lose their energy to radiation and conditions in the flow change. Ultimately, the particles that gave rise to γ -ray radiation near the star emit radio far from the binary. The flow changes from a cometary shape to a spiral-shaped nebula with a step $\sim \sigma c P_{\text{orb}}$. Matching with the observed size of radio nebula in LS 5039 and LSI +61°303 implies $\sigma \approx 0.01$. On large scales (several turns of the spiral), the radio emission is underestimated by the model. Changes in the flow conditions on such scales, notably velocity and magnetic field evolution, are undoubtedly more complex than the simple assumptions made here. Nevertheless, the combination of the peculiar geometry, emission at various distances and inclination can conspire to create radio maps with seemingly bipolar ejections of material as in X-ray binary jets. Predictions are made for LS 5039 that are amenable to observation with radio interferometry.

The present model does not provide an entirely satisfactory view of the GeV γ -rays detected by EGRET in LS 5039 and LSI +61°303. The fluxes are in both cases underproduced. The discrepancy might have an observational side, with fluxes overestimated, comprising the combined emission from several sources in the wide instrumental PSF. Much more precise measurements will be made by *GLAST* in the future. On the theoretical side, inadequacies in the treatment of the emission region might be at work, notably anisotropies. The stellar radiation is anisotropic, the pulsar wind is anisotropic, the termination shock is not spherical etc. If synchrotron losses are limited, IC cooling of particles will enhance the GeV yield and lower the discrepancy. The mismatch between EGRET and HESS spectra of LS 5039 is also difficult for leptonic jet models (Dermer & Böttcher, 2006; Paredes et al., 2006).

The present model does not either satisfyingly explain the nature of the radio outbursts in LSI +61°303 and PSR B1259-63. However, the scenario provides a unifying context. As has been argued for both sources, the reason must be found in the Be nature of the companion stars. The progression of the pulsar in the slow, dense wind could lead to an enclosed geometry, rather than an open comet tail geometry. Radio emission might also be enhanced by the acceleration of particles from the stellar wind itself. There is as yet no coherent picture of the radio outbursts of LSI +61°303 in the context of the accretion-driven / relativistic jet scenario.

Despite these shortcomings, the explanatory scope of the model is large. Once the orbital parameters are set, the only variables that come into play are \dot{E} , σ , γ_p and R_s . The SEDs come out naturally for reasonable inputs and the comet-tail nebula produces radio emission on scales of $\sigma P_{\text{orb}} c$ and beyond. The pulsar scenario therefore seems like a viable alternative to the accretion/jet scenario for LS 5039 and LSI +61°303.

Acknowledgements. I thank B. Giebels, J.-P. Lasota, M. Sikora, R. Taam and R. Fender for their comments on this work.

References

- Aharonian, F. A., Anchordoqui, L. A., Khangulyan, D., & Montaruli, T. 2006, *Journal Phys.*, in press (astro-ph/0508658)
- Aharonian, F. A., et al. (HESS collaboration) 2005a, *Science*, 309, 746
- Aharonian, F. A., et al. (HESS collaboration) 2005b, *A&A*, 442, 1
- Arons, J., & Tavani, M. 1993, *ApJ*, 403, 249
- Arons, J., & Tavani, M. 1994, *ApJS*, 90, 797
- Bahcall, J. N. 1978, *ARA&A*, 16, 241
- Ball, L., & Kirk, J. G. 2000, *Astroparticle Physics*, 12, 335
- Ball, L., Melatos, A., Johnston, S., & Skjaeraasen, O., 1999, *ApJ*, 514, L39
- Bednarek, W., 2006, *MNRAS*, 368, 579
- Blumenthal G. R., & Gould R. J., 1970, *Rev. Mod. Phys.*, 42, 237
- Bogovalov, S. V., & Aharonian, F. A. 2000, *MNRAS*, 313, 504
- Bosch-Ramon, V., Paredes, J. M., Ribó, M., Miller, J. M., Reig, P., & Martí, J. 2005, *ApJ*, 628, 388
- Böttcher, M., & Dermer, C. D. 2005, *ApJ*, 634, L81
- Bucciantini, N., Amato, E., & Del Zanna, L. 2005, *A&A*, 434, 189
- Casares, J., Ribas, I., Paredes, J. M., Martí, J., & Allende Prieto, C. 2005a, *MNRAS*, 360, 1105
- Casares, J., Ribó, M., Ribas, I., Paredes, J. M., Martí, J., & Herrero, A. 2005b, *MNRAS*, 364, 899
- Chang, J. S., & Cooper, G., 1970, *J. Comp. Phys.*, 6, 1
- Clark, J. S., et al. 2001, *A&A*, 376, 476
- Connors, T. W., Johnston, S., Manchester, R. N., & McConnell, D., 2002, *MNRAS*, 336, 1201
- Conti, P. S. 1978, *A&A*, 63, 225
- Cutri, R. M., et al. 2003, *VizieR Online Data Catalog*, 2246, 0
- Dermer, C. D., & Böttcher, M. 2006, *ApJ*, in press (astro-ph/0512162)
- Dougherty, S. M., Pittard, J. M., Kasian, L., Coker, R. F., Williams, P. M., & Lloyd, H. M. 2003, *A&A*, 409, 217
- Dubus, G., 2006, *A&A*, 451, 9
- Fegan, S. J., et al. 2005, *ApJ*, 624, 638
- Fender, R. P., Belloni, T. M., & Gallo, E. 2004, *MNRAS*, 355, 1105
- Fender, R. P., 2006, chapter 9 in 'Compact Stellar X-Ray Sources', eds. W.H.G. Lewin and M. van der Klis, Cambridge Astrophysics Series 39, Cambridge University Press, pp381-420
- Frail, D. A., Seaquist, E. R., & Taylor, A. R. 1987, *AJ*, 93, 1506
- Gaensler, B. M. 2005, *Advances in Space Research*, 35, 1116
- Gaensler, B. M., & Slane, P. O. 2006, to appear in volume 44 of *ARA&A* (astro-ph/0601081)
- Gould R. J., & Schröder G. P., 1967, *Phys. Rev.*, 155, 1404
- Gregory, P. C. 2002, *ApJ*, 575, 427
- Gregory, P. C., et al. 1979, *AJ*, 84, 1030
- Gregory, P. C., & Taylor, A. R. 1978, *Nature*, 272, 704
- Greiner, J., & Rau, A. 2001, *A&A*, 375, 145
- Grove, J. E., Tavani, M., Purcell, W. R., Johnson, W. N., Kurfess, J. D., Strickman, M. S., & Arons, J. 1995, *ApJ*, 447, L113
- Hall, T. A. et al. 2003, *ApJ*, 583, 853
- Harmon, B. A., et al. 2004, *ApJS*, 154, 585
- Harrison, F. A., Ray, P. S., Leahy, D. A., Waltman, E. B., & Pooley, G. G. 2000, *ApJ*, 528, 454
- Hartman, R. C., et al. 1999, *ApJS*, 123, 79
- Hoffman, C. M., Sinnis, C., Fleury, P., & Punch, M. 1999, *Reviews of Modern Physics*, 71, 897
- Hutchings, J. B., & Crampton, D. 1981, *PASP*, 93, 486
- Illarionov, A. F., & Sunyaev, R. A. 1975, *A&A*, 39, 185
- Johnston, S., Manchester, R. N., Lyne, A. G., Bailes, M., Kaspi, V. M., Qiao, G., & D'Amico, N. 1992, *ApJ*, 387, L37
- Johnston, S., Manchester, R. N., Lyne, A. G., Nicastro, L., & Spyromilio, J. 1994, *MNRAS*, 268, 430
- Johnston, S., Ball, L., Wang, N., & Manchester, R. N. 2005, *MNRAS*, 358, 1069
- Jones F. C., 1968, *Phys. Rev.*, 167, 1159
- Kennel, C. F., & Coroniti, F. V. 1984a, *ApJ*, 283, 694
- Kennel, C. F., & Coroniti, F. V. 1984b, *ApJ*, 283, 710
- Kirk, J. G., Ball, L., & Skjaeraasen, O. 1999, *Astroparticle Physics*, 10, 31
- Leahy, D. A. 2004, *A&A*, 413, 1019
- Leahy, D. A., Harrison, F. A., & Yoshida, A. 1997, *ApJ*, 475, 823
- Manchester, R. N., Johnston, S., Lyne, A. G., D'Amico, N., Bailes, M., & Nicastro, L. 1995, *ApJ*, 445, L137
- Maraschi, L., & Treves, A. 1981, *MNRAS*, 194, 1P
- Maraschi, L., Treves, A., & Tanzi, E. G. 1981, *ApJ*, 248, 1010
- Marraco, L. G., & Orsatti, A. M. 1982, *Revista Mexicana de Astronomia y Astrofisica*, 5, 183
- Martí, J., & Paredes, J. M. 1995, *A&A*, 298, 151
- Martí, J., Peracaula, M., Paredes, J. M., Massi, M., & Estalella, R. 1998a, *A&A*, 329, 951
- Martí, J., Paredes, J. M., & Ribó, M. 1998b, *A&A*, 338, L71
- Martocchia, A., Motch, C., & Negueruela, I. 2005, *A&A*, 430, 245
- Massi, M., Paredes, J. M., Estalella, R., & Felli, M. 1993, *A&A*, 269, 249
- Massi, M., Ribó, M., Paredes, J. M., Peracaula, M., & Estalella, R. 2001, *A&A*, 376, 217
- Massi, M., Ribó, M., Paredes, J. M., Garrington, S. T., Peracaula, M., & Martí, J. 2004, *A&A*, 414, L1
- McSwain, M. V., & Gies, D. R. 2002, *ApJ*, 568, L27
- McSwain, M. V., Gies, D. R., Huang, W., Wiita, P. J., Wingert, D. W., & Kaper, L. 2004, *ApJ*, 600, 927
- Melatos, A., Johnston, S., & Melrose, D. B. 1995, *MNRAS*, 275, 381
- Meurs, E. J. A., & van den Heuvel, E. P. J. 1989, *A&A*, 226, 88
- Moderski, R., Sikora, M., Coppi, P. S., & Aharonian, F. 2005, *MNRAS*, 355, 845
- Motch, C., Haberl, F., Dennerl, K., Pakull, M., & Janot-Pacheco, E. 1997, *A&A*, 323, 853
- Murata, K., Tamaki, H., Maki, H., & Shibazaki, N., 2003, *PASJ*, 55, 473
- Paredes, J. M., Martí, J., Estalella, R., & Sarrate, J. 1991, *A&A*, 248, 124
- Paredes, J. M., Martí, J., Peracaula, M., & Ribó, M. 1997, *A&A*, 320, L25

- Paredes, J. M., Martí, J., Ribó, M., & Massi, M. 2000, *Science*, 288, 2340
- Paredes, J. M., Ribó, M., Ros, E., Martí, J., & Massi, M. 2002, *A&A*, 393, L99
- Paredes, J. M., Bosch-Ramon, V., & Romero, G. E. 2006, *A&A*, 451, 259
- Rees, M. J., & Gunn, J. E. 1974, *MNRAS*, 167, 1
- Reig, P., Ribó, M., Paredes, J. M., & Martí, J. 2003, *A&A*, 405, 285
- Ribó, M., Reig, P., Martí, J., & Paredes, J. M. 1999, *A&A*, 347, 518
- Ribó, M., Paredes, J. M., Romero, G. E., Benaglia, P., Martí, J., Fors, O., & García-Sánchez, J. 2002, *A&A*, 384, 954
- Romero, G. E., Christiansen, H. R., & Orellana, M. 2005, *ApJ*, 632, 1093
- Stirling, A. M., Spencer, R. E., de la Force, C. J., Garrett, M. A., Fender, R. P., & Ogle, R. N. 2001, *MNRAS*, 327, 1273
- Strickman, M. S., Tavani, M., Coe, M. J., Steele, I. A., Fabregat, J., Martí, J., Paredes, J. M., & Ray, P. S. 1998, *ApJ*, 497, 419
- Tavani M., & Arons J., 1997, *ApJ*, 477, 439
- Tavani, M., Arons, J., & Kaspi, V. M. 1994, *ApJ*, 433, L37
- Tavani, M., et al. 1996, *A&AS*, 120, 221
- Taylor, A. R., & Gregory, P. C. 1982, *ApJ*, 255, 210
- Taylor, A. R., & Gregory, P. C. 1984, *ApJ*, 283, 273
- van der Klis, M. 2006, chapter 2 in 'Compact Stellar X-Ray Sources', eds. W.H.G. Lewin and M. van der Klis, Cambridge Astrophysics Series 39, Cambridge University Press, pp39-112
- Wang, Q. D., Li, Z.-Y., & Begelman, M. C. 1993, *Nature*, 364, 127
- Waters, L. B. F. M. 1986, *A&A*, 162, 121
- Waters, L. B. F. M., van den Heuvel, E. P. J., Taylor, A. R., Habets, G. M. H. J., & Persi, P. 1988, *A&A*, 198, 200
- Zahn, J.-P. 1977, *A&A*, 57, 383




Fig. 1. Observed spectral energy distributions of LS 5039, LSI +61°303 and PSR B1259-63. Fluxes have been transformed to spectral luminosities assuming distances of 2.5 kpc (LS 5039, Casares et al. 2005b), 2.3 kpc (LSI +61°303, Gregory et al. 1979) and 1.5 kpc (PSR B1259-63, Johnston et al. 1994). The optical-UV emission peaking at ~ 1 eV in the binaries is the light from the stellar companion (uncorrected for absorption by the interstellar medium). Radio to X-ray flux values in LS 5039 are taken from (in order of increasing frequency) Martí et al. (1998b); Clark et al. (2001); Martocchia et al. (2005); in LSI +61°303 from Strickman et al. (1998); Martí & Paredes (1995); Maraschi et al. (1981); Leahy et al. (1997); Greiner & Rau (2001); in PSR B1259-63 from Johnston et al. (2005); Cutri et al. (2003); Marraco & Orsatti (1982). The *RXTE* spectrum of PSR B1259-63 obtained during the 2004 periastron passage was kindly provided by B. Giebels. The upper envelope of radio measurements in LSI +61°303 and PSR B1259-63 corresponds to values during radio outbursts. The hard X-ray BATSE points in LS 5039 and LSI +61°303 are from Harmon et al. (2004); the OSSE bowtie is from Grove et al. (1995) and the HE γ -ray EGRET boxes are from Hartman et al. (1999); EGRET upper limits in PSR B1259-63 are from Tavani et al. (1996). The VHE γ -ray HESS measurements are from Aharonian et al. (2005a,b) and the Whipple upper limits for LSI +61°303 from Fegan et al. (2005).




Fig. 2. Timescales at the standoff point as a function of the electron Lorentz factor γ in LS 5039. The radiative timescale $t_{\text{rad}}^{-1} = t_{\text{sync}}^{-1} + t_{\text{IC}}^{-1}$ (solid lines) and the escape timescale t_{esc} (dashed lines) are shown at periastron (black) and apastron (grey). Synchrotron cooling dominates above $\gamma_S \approx 2 \cdot 10^6$. Inverse Compton cooling dominates below, the transition between Thomson and Klein-Nishina regimes occurring at $\gamma_{\text{KN}} \approx 6 \cdot 10^4$. R_s is $2 \cdot 10^{11}$ cm at periastron ($d = 0.1$ AU) and $4 \cdot 10^{11}$ cm at apastron ($d = 0.2$ AU) (see Table 2).




Fig. 5. Large scale appearance of the pulsar nebula around LS 5039 at periastron. In practice, limited angular resolution smoothes out the emission from what is shown here (see Fig. 7). The binary is located at the origin. Its size of a few tenths of AU is much smaller than the scale of the map. Material originating from the standoff point travels outwards following the direction of the comet tail at each orbital phase (shown by grey lines; the projected outflow path at apastron is indicated by a dark line). Because of the orbital motion, the emission winds up into a spiral. To illustrate this, dots are plotted at the locations reached by material expelled at *regular* time intervals along the orbit. The total elapsed timescale represented corresponds to almost an orbital period and a half. The greyscale codes the intensity of the 5 GHz radio emission (square root scale). At large distances, the step of the spiral tends to $\sigma c P_{\text{orb}}$ which is ≈ 7 AU for the adopted parameters (same as those of Figs. 3-4). The inclination ($i = 120^\circ$, corresponding to a retrograde pulsar orbit, see §6.1.2), and the major axis orientation ($\omega = 226^\circ$) used here are consistent with the measurements of the companion star's radial velocity (see orbit schematic in Fig. 6 of Casares et al. 2005b).




Fig. 6. Spectral energy distribution of LS 5039 at periastron (dark full line) and apastron (dark dashed line). The corresponding grey lines show the fraction received by the observer after absorption of the VHE γ -rays on stellar photons. Parameters are identical to those of Figs. 3-4 (but using $R_s = 4 \cdot 10^{11}$ cm at apastron). Changes in the overall SED are limited and only minor variability is expected except at VHE energies where $\gamma\gamma$ absorption strongly modulates the flux on the orbital period.




Fig. 9. Left: Map of the 5 GHz radio emission from the pulsar nebula in LSI +61°303 at $\phi = 0.5$ ($i = 120^\circ$). ϕ is zero at periastron passage (for which $\phi_{\text{radio}} = 0.23$, Gregory 2002). Right: Corresponding outflow path (as in Fig. 5 but on a linear greyscale) before smoothing (the projected outflow path at apastron is indicated by a dark line). The stellar wind is assumed to be polar (top panel of Fig. 8). The smoothing Gaussian kernel has a FWHM of 8 mas ($2.8 \cdot 10^{14}$ cm at 2.3 kpc). Contours are at 50%, 25% and 10% of maximum emission. There is a preferred direction to the right, towards apastron.




Fig. 8. Spectral energy distribution of LSI +61°303 at periastron (full black lines) and apastron (dashed black lines). The pulsar is assumed to have $\dot{E} = 10^{36}$ erg s $^{-1}$ and $\gamma = 10^5$ as for LS 5039, but a slightly higher $\sigma = 0.02$. Top: The stellar wind is polar with a standoff point at $R_s = 8 \cdot 10^{11}$ cm (periastron) and $4 \cdot 10^{12}$ cm (apastron). Bottom: The stellar wind is equatorial with $R_s = 5 \cdot 10^{10}$ cm (periastron) and $2 \cdot 10^{12}$ cm (apastron). Grey lines show the fraction of the emission occurring when $\Delta t > P_{\text{orb}}$ (the periastron line is visible in the top panel but not in the bottom panel).




Fig. 10. Spectral energy distribution of PSR B1259-63 at periastron (solid line) and apastron (dashed line). The pulsar is assumed to have $\dot{E} = 8 \cdot 10^{35}$ erg s $^{-1}$, $\gamma = 10^5$ and $\sigma = 0.01$. The stellar wind is polar (same parameters as LSI +61°303) with a standoff point at $R_s = 3 \cdot 10^{12}$ cm (periastron) and $4 \cdot 10^{13}$ cm (apastron). With an equatorial, slow wind, the standoff point at periastron is at $R_s = 10^{12}$ cm, not different enough from the polar wind case to significantly change the expected SED from that shown above. The grey dashed line shows the summed late time emission ($\Delta t > P_{\text{orb}}$) at apastron, which occurs on scales $\gtrsim 0.01$ pc.




Fig. 11. Left: Map of 5 GHz radio emission from PSR B1259-63 close to periastron passage ($\phi = 0.1$, $i = 30^\circ$), assuming only a polar-type stellar wind. Right: Corresponding outflow path on the same scale before smoothing (the projected outflow path at apastron is indicated by a dark line). The smoothing gaussian kernel has a FWHM of $0.05''$ ($1.1 \cdot 10^{15}$ cm at 1.5 kpc). The contours are at 50%, 25% and 10% of the maximum intensity. The tail extends to a few arcseconds. The spatial scale is large given the size of the orbit (hence R_s); with an even larger eccentricity than LSI +61°303, the preferred direction of the nebula towards apastron is very clear.